SANTA CRUZ BIOTECHNOLOGY, INC.

rhodopsin (C7): sc-53991



BACKGROUND

G protein-coupled receptors (GPCRs), which are characterized as containing seven transmembrane α helices, elicit G protein-mediated signaling cascades in response to a variety of stimuli. The opsin subfamily, which represents approximately 90 percent of all GPCRs, is comprised of photoreceptors that are activated by light, which include the red, green and blue-sensitive opsins and rhodopsin. The opsin subfamily consists of an apoprotein covalently linked to 11-*cis*-retinal, which undergoes isomerization upon the absorption of photons. This isomerization leads to a conformational change of the protein which results in the activation of hundreds of G proteins. Specifically, rhodopsin exhibits a maximal absorption at 495 nm and mediates vision in dim light. Mutations in the rhodopsin gene lead to retinitis pigmentosa, which can be inherited as an autosomal dominant, an autosomal recessive or an X-linked recessive disorder.

REFERENCES

- Fung, B.K., et al. 1981. Flow of information in the light-triggered cyclic nucleotide cascade of vision. Proc. Natl. Acad. Sci. USA 78: 152-156.
- 2. Hargrave, P.A., et al. 1983. The structure of bovine rhodopsin. Biophys. Struct. Mech. 9: 235-244.
- Wang, S.Z., et al. 1992. A visual pigment from chicken that resembles rhodopsin: amino acid sequence, gene structure and functional expression. Biochemistry 13: 3309-3315.
- al-Maghtheh, M., et al. 1993. Rhodopsin mutations in autosomal dominant retinitis pigmentosa. Hum. Mutat. 2: 249-255.
- 5. liri, T., et al. 1998. G protein diseases furnish a model for the turn-on switch. Nature 394: 35-38.
- Lindsay, S.M., et al. 1999. Spectral sensitivity of vision and bioluminescence in the midwater shrimp. Biol. Bull. 197: 348-360.
- 7. Palczewski, K., et al. 2000. Crystal structure of rhodopsin: a G proteincoupled receptor. Science 289: 739-745.

CHROMOSOMAL LOCATION

Genetic locus: Rho (mouse) mapping to 6 E3.

SOURCE

rhodopsin (C7) is a mouse monoclonal antibody raised against rhodopsin of bovine origin.

PRODUCT

Each vial contains 200 μg IgG_1 kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, **D0 NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

rhodopsin (C7) is recommended for detection of rhodopsin of mouse, rat and bovine origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)] and immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

Suitable for use as control antibody for rhodopsin siRNA (m): sc-40151, rhodopsin shRNA Plasmid (m): sc-40151-SH and rhodopsin shRNA (m) Lentiviral Particles: sc-40151-V.

Molecular Weight of rhodopsin: 40 kDa.

Positive Controls: CTLL-2 cell lysate: sc-2242, mouse eye extract: sc-364241 or rat eye extract: sc-364805.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz MarkerTM Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

DATA





rhodopsin (C7): sc-53991. Western blot analysis of rhodopsin expression in mouse eye tissue extract (A) and CTLL-2 whole cell lysate (B). rhodopsin (C7): sc-53991. Western blot analysis of rhodopsin expression in mouse eye (A) and rat eye (B) tissue extract.

SELECT PRODUCT CITATIONS

- Roman, D., et al. 2016. Ocular toxicity of AUY922 in pigmented and albino rats. Toxicol. Appl. Pharmacol. 309: 55-62.
- Shamsnajafabadi, H., et al. 2022. Neural differentiation of human retinal pigment epithelial cells on alginate/gelatin substrate. Mol. Vis. 28: 412-431.



See **rhodopsin (1D4): sc-57432** for rhodopsin antibody conjugates, including AC, HRP, FITC, PE, and Alexa Fluor[®] 488, 546, 594, 647, 680 and 790.