

COX10 (E-17): sc-54062

BACKGROUND

Cytochrome c oxidase (COX) localizes to the mitochondrial inner membrane and is the terminal enzyme in the electron transfer chain, functioning as a transmembrane proton pump that builds an electrochemical gradient with chemical energy from the reduction of O₂. The COX subunit 10, or COX10 (also known as heme A:farnesyltransferase or heme O synthase), is a multi-pass transmembrane protein encoded by a nuclear gene. COX10 was originally identified in yeast and its structure is conserved from *E. coli* to human. COX10 is responsible for catalyzing the first step in the biosynthesis of heme A: the conversion of protoheme (heme B) to heme O by the addition of a farnesyl group. As a result, COX10 is necessary for the expression of a functional COX. A mutation in the gene encoding COX10 may result in COX deficiency in humans.

REFERENCES

- Glerum, D.M. and Tzagoloff, A. 1994. Isolation of a human cDNA for heme A:farnesyltransferase by functional complementation of a yeast COX10 mutant. *Proc. Natl. Acad. Sci. USA* 91: 8452-8456.
- Murakami, T., Reiter, L.T. and Lupski, J.R. 1997. Genomic structure and expression of the human heme A:farnesyltransferase (COX10) gene. *Genomics* 42: 161-164.
- Valnot, I., von Kleist-Retzow, J.C., Barrientos, A., Gorbatyuk, M., Taanman, J.W., Mehaye, B., Rustin, P., Tzagoloff, A., Munnich, A. and Rötig, A. 2000. A mutation in the human heme A:farnesyltransferase gene (COX10) causes cytochrome c oxidase deficiency. *Hum. Mol. Genet.* 9: 1245-1249.
- Antonicka, H., Leary, S.C., Guercin, G.H., Agar, J.N., Horvath, R., Kennaway, N.G., Harding, C.O., Jaksch, M. and Shoubbridge, E.A. 2003. Mutations in COX10 result in a defect in mitochondrial heme A biosynthesis and account for multiple, early-onset clinical phenotypes associated with isolated COX deficiency. *Hum. Mol. Genet.* 12: 2693-2702.
- Coenen, M.J., van den Heuvel, L.P., Ugalde, C., Ten Brinke, M., Nijtmans, L.G., Trijbels, F.J., Beblo, S., Maier, E.M., Muntau, A.C. and Smeitink, J.A. 2004. Cytochrome c oxidase biogenesis in a patient with a mutation in COX10 gene. *Ann. Neurol.* 56: 560-564.
- Diaz, F., Thomas, C.K., Garcia, S., Hernandez, D. and Moraes, C.T. 2005. Mice lacking COX10 in skeletal muscle recapitulate the phenotype of progressive mitochondrial myopathies associated with cytochrome c oxidase deficiency. *Hum. Mol. Genet.* 14: 2737-2748.

CHROMOSOMAL LOCATION

Genetic locus: COX10 (human) mapping to 17p12; Cox10 (mouse) mapping to 11 B3.

SOURCE

COX10 (E-17) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of COX10 of human origin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-54062 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

COX10 (E-17) is recommended for detection of COX10 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

COX10 (E-17) is also recommended for detection of COX10 in additional species, including equine, canine, bovine, porcine and avian.

Suitable for use as control antibody for COX10 siRNA (h): sc-72303, COX10 siRNA (m): sc-72304, COX10 shRNA Plasmid (h): sc-72303-SH, COX10 shRNA Plasmid (m): sc-72304-SH, COX10 shRNA (h) Lentiviral Particles: sc-72303-V and COX10 shRNA (m) Lentiviral Particles: sc-72304-V.


Molecular Weight of COX10: 51 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200, HEK293 whole cell lysate: sc-45136 or CCRF-CEM cell lysate: sc-2225.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



COX10 (E-17): sc-54062. Western blot analysis of COX10 expression in HeLa (A), HEK293 (B) and CCRF-CEM (C) whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.