

# SID-2 (G-17): sc-54150

## BACKGROUND

SID-1 and SID-2 belong to the systemic RNA interference defective-1 (SID1) family of transmembrane proteins. SID-1, originally identified in *C. elegans*, is an 827 amino acid long protein. It localizes to the cell membrane and contains 11 transmembrane domains. This suggests that SID-1 possibly functions as a channel protein. The overexpression of SID-1 enhances double stranded RNA (dsRNA) uptake in pancreatic ductal adenocarcinoma cells. SID-2, also first identified in *C. elegans*, is an 832 amino acid long protein with multiple transmembrane domains. At least two isoforms exist for SID-2 due to alternative splicing. Isoform 2 contains an additional 21 amino acids after residue 387 and has an alternate sequence that is 8 amino acids shorter for residues 814 to 832 of isoform 1.

## REFERENCES

1. Winston, W.M., Molodowitch, C. and Hunter, C.P. 2002. Systemic RNAi in *C. elegans* requires the putative transmembrane protein SID-1. *Science* 295: 2456-2459.
2. Feinberg, E.H. and Hunter, C.P. 2003. Transport of dsRNA into cells by the transmembrane protein SID-1. *Science* 301: 1545-1547.
3. Duxbury, M.S., Ashley, S.W. and Whang, E.E. 2005. RNA interference: a mammalian SID-1 homologue enhances siRNA uptake and gene silencing efficacy in human cells. *Biochem. Biophys. Res. Commun.* 331: 459-463.
4. Kim, J.K., Gabel, H.W., Kamath, R.S., Tewari, M., Pasquinelli, A., Rual, J.F., Kennedy, S., Dybbs, M., Bertin, N., Kaplan, J.M., Vidal, M. and Ruvkun, G. 2005. Functional genomic analysis of RNA interference in *C. elegans*. *Science* 308: 1164-1167.
5. Tokue, I., Yamasaki, K. and Nanbu, S. 2005. Vibrational energies for the X1A1, A1B1, and B1A1 states of SiH2/SID-2 and related transition probabilities based on global potential energy surfaces. *J. Chem. Phys.* 122: 144307.
6. Tsang, S.Y., Moore, J.C., Huizen, R.V., Chan, C.W. and Li, R.A. 2007. Ectopic expression of systemic RNA interference defective protein in embryonic stem cells. *Biochem. Biophys. Res. Commun.* 357: 480-486.
7. Winston, W.M., Sutherlin, M., Wright, A.J., Feinberg, E.H. and Hunter, C.P. 2007. *Caenorhabditis elegans* SID-2 is required for environmental RNA interference. *Proc. Natl. Acad. Sci. USA* 104: 10565-10570.
8. Hunter, C.P., Winston, W.M., Molodowitch, C., Feinberg, E.H., Shih, J., Sutherlin, M., Wright, A.J. and Fitzgerald, M.C. 2007. Systemic RNAi in *Caenorhabditis elegans*. *Cold Spring Harb. Symp. Quant. Biol.* 71: 95-100.

## CHROMOSOMAL LOCATION

Genetic locus: SIDT2 (human) mapping to 11q23.3; Sidt2 (mouse) mapping to 9 A5.2.

## SOURCE

SID-2 (G-17) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an extracellular domain of SID-2 of human origin.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-54150 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## APPLICATIONS

SID-2 (G-17) is recommended for detection of SID-2 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

SID-2 (G-17) is also recommended for detection of SID-2 in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for SID-2 siRNA (h): sc-72282, SID-2 siRNA (m): sc-72283, SID-2 shRNA Plasmid (h): sc-72282-SH, SID-2 shRNA Plasmid (m): sc-72283-SH, SID-2 shRNA (h) Lentiviral Particles: sc-72282-V and SID-2 shRNA (m) Lentiviral Particles: sc-72283-V.

Molecular Weight of SID-2: 104 kDa.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.