

AMID (S-20): sc-54163

BACKGROUND

AMID (apoptosis-inducing factor (AIF)-like mitochondrion-associated inducer of death), also called p53-responsive gene 3 (PRG3), is a member of the FAD-dependent oxidoreductase family. AMID is a caspase-independent pro-apoptotic flavoprotein with NAD(P)H oxidase activity localizing to the cytosol and associated with the outer mitochondrial membrane. AMID shares significant homology with AIF and NADH-oxidoreductases. It is expressed in most normal tissues and its expression is upregulated by p53. Two AMID isoforms exist due to alternative splicing. Isoform 1 is the full length protein and isoform 2 is missing amino acids 99-138. Isoform 2 also has an additional three amino acids inserted after residue 206. Overexpression of AMID leads to apoptosis.

REFERENCES

1. Wu, M., Xu, L.G., Li, X., Zhai, Z. and Shu, H.B. 2002. AMID, an apoptosis-inducing factor-homologous mitochondrion-associated protein, induces caspase-independent apoptosis. *J. Biol. Chem.* 277: 25617-25623.
2. Lü, C.X., Fan, T.J., Hu, G.B. and Cong, R.S. 2003. Apoptosis-inducing factor and apoptosis. *Sheng Wu Hua Xue Yu Sheng Wu Wu Li Xue Bao* 35: 881-885.
3. Jan, Y., Matter, M., Pai, J.T., Chen, Y.L., Pilch, J., Komatsu, M., Ong, E., Fukuda, M. and Ruoslahti, E. 2004. A mitochondrial protein, Bit1, mediates apoptosis regulated by integrins and Groucho/TLE corepressors. *Cell* 116: 751-762.
4. Wu, M., Xu, L.G., Su, T., Tian, Y., Zhai, Z. and Shu, H.B. 2004. AMID is a p53-inducible gene downregulated in tumors. *Oncogene* 23: 6815-6819.
5. Marshall, K.R., Gong, M., Wodke, L., Lamb, J.H., Jones, D.J., Farmer, P.B., Scrutton, N.S. and Munro, A.W. 2005. The human apoptosis-inducing protein AMID is an oxidoreductase with a modified flavin cofactor and DNA binding activity. *J. Biol. Chem.* 280: 30735-30740.
6. Hu, Q., Chang, J., Tao, L., Yan, G., Xie, M. and Wang, Z. 2005. Endoplasmic reticulum mediated necrosis-like apoptosis of HeLa cells induced by Ca²⁺ oscillation. *J. Biochem. Mol. Biol.* 38: 709-716.
7. Li, W., Sun, L., Liang, Q., Wang, J., Mo, W. and Zhou, B. 2006. Yeast AMID homologue Ndi1p displays respiration-restricted apoptotic activity and is involved in chronological aging. *Mol. Biol. Cell* 17: 1802-1811.
8. Mei, J., Webb, S., Zhang, B. and Shu, H.B. 2006. The p53-inducible apoptotic protein AMID is not required for normal development and tumor suppression. *Oncogene* 25: 849-856.
9. Varecha, M., Amrichová, J., Zimmermann, M., Ullman, V., Lukášová, E. and Kozubek, M. 2007. Bioinformatic and image analyses of the cellular localization of the apoptotic proteins endonuclease G, AIF, and AMID during apoptosis in human cells. *Apoptosis* 12: 1155-1171.

CHROMOSOMAL LOCATION

Genetic locus: AIFM2 (human) mapping to 10q22.1; Aifm2 (mouse) mapping to 10 B4.

SOURCE

AMID (S-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of AMID of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-54163 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

AMID (S-20) is recommended for detection of AIF-like mitochondrion-associated inducer of death of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

AMID (S-20) is also recommended for detection of AIF-like mitochondrion-associated inducer of death in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for AMID siRNA (h): sc-72339, AMID siRNA (m): sc-72340, AMID shRNA Plasmid (h): sc-72339-SH, AMID shRNA Plasmid (m): sc-72340-SH, AMID shRNA (h) Lentiviral Particles: sc-72339-V and AMID shRNA (m) Lentiviral Particles: sc-72340-V.

Molecular Weight of AMID: 41 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.