



Pancreatic Icosapeptide (M-15): sc-54291

BACKGROUND

Pancreatic Polypeptide (PP) and Pancreatic Icosapeptide (PI) are both deduced from the pancreatic prohormone precursor. The two peptide sequences are separated by a Gly-Lys-Arg cleavage and amidation site. The Pancreatic Polypeptide lies on the N-terminal side of this cleavage site while the Pancreatic Icosapeptide lies on the C-terminal side. The prohormone precursor is produced by the endocrine F cells of the pancreatic islets and, in response to food intake, the Pancreatic Polypeptide is released into the circulation. The Pancreatic Polypeptide is a member of the neuropeptide Y (NPY) family of hormones that bind to Y receptors. In particular, it binds to Y4 receptors and functions as an anorexigenic hormone. Subjects with Prader-Willi syndrome have a decreased Pancreatic Polypeptide response to a meal. Administration of Pancreatic Polypeptide decreases food intake and may serve as a therapeutic option for treatment of obesity.

REFERENCES

- Boel, E., Schwartz, T.W., Norris, K.E. and Fiil, N.P. 1984. A cDNA encoding a small common precursor for human Pancreatic Polypeptide and Pancreatic Icosapeptide. *EMBO J.* 3: 909-912.
- Schwartz, T.W., Hansen, H.F., Håkanson, R., Sundler, F. and Tager, H.S. 1984. Human Pancreatic Icosapeptide: isolation, sequence, and immunocytochemical localization of the COOH-terminal fragment of the Pancreatic Polypeptide precursor. *Proc. Natl. Acad. Sci. US* 81: 708-712.
- Leiter, A.B., Montminy, M.R., Jamieson, E. and Goodman, R.H. 1985. Exons of the human Pancreatic Polypeptide gene define functional domains of the precursor. *J. Biol. Chem.* 260: 13013-13017.
- Takeuchi, T., Gumucio, D.L., Yamada, T., Meisler, M.H., Minth, C.D., Dixon, J.E., Eddy, R.E. and Shows, T.B. 1986. Genes encoding Pancreatic Polypeptide and neuropeptide Y are on human chromosomes 17 and 7. *J. Clin. Invest.* 77: 1038-1041.
- Koska, J., DelParigi, A., de Courten, B., Weyer, C. and Tataranni, P.A. 2004. Pancreatic Polypeptide is involved in the regulation of body weight in Pima Indian male subjects. *Diabetes* 53: 3091-3096.
- Schmidt, P.T., Näslund, E., Grybäck, P., Jacobsson, H., Holst, J.J., Hilsted, L. and Hellström, P.M. 2005. A role for Pancreatic Polypeptide in the regulation of gastric emptying and short-term metabolic control. *J. Clin. Endocrinol. Metab.* 90: 5241-5246.
- Ku, S.K., Lee, H.S. and Lee, J.H. 2006. The regional distribution and relative frequency of gastrointestinal endocrine cells in the nude mice, Balb/c-nu/nu: an immunohistochemical study. *Anat. Histol. Embryol.* 35: 104-110.
- Kapran, Y., Bauersfeld, J., Anlauf, M., Sipos, B. and Klöppel, G. 2006. Multihormonality and entrapment of islets in pancreatic endocrine tumors. *Virchows Arch.* 448: 394-398.
- Yamada, S., Terada, K., Ueno, Y., Sugiyama, T., Seno, M. and Kojima, I. 2006. Differentiation of adult hepatic stem-like cells into pancreatic endocrine cells. *Cell Transplant.* 14: 647-653.

CHROMOSOMAL LOCATION

Genetic locus: Ppy (mouse) mapping to 11 D.

SOURCE

Pancreatic Icosapeptide (M-15) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of pancreatic prohormone precursor of mouse origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-54291 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

Pancreatic Icosapeptide (M-15) is recommended for detection of Pancreatic Icosapeptide and pancreatic prohormone precursor of mouse origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for Pancreatic Polypeptide siRNA (m): sc-62750.

Molecular Weight of Pancreatic Icosapeptide precursor: 12 kDa.

Molecular Weight of Pancreatic Icosapeptide: 2 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.