

SUN1 (N-19): sc-54447

BACKGROUND

UNC84A (UNC84 homolog A), also known as SUN1, is a multi-pass nuclear membrane protein that is involved in nuclear anchoring and migration. Highly expressed in heart, brain and testis, UNC84A functions as an A-type Lamin-binding protein that forms a link between the inner and outer nuclear envelope membranes. This link acts as a structural bridge between the nuclear interior and the Actin cytoskeleton and is essential for proper localization of nuclear envelope proteins. Additionally, UNC84A may be involved in telomere attachment and in normal testis development. UNC84A contains one UNC84 (SUN) domain and exists as four isoforms due to alternative splicing events.

REFERENCES

1. Dreger, M., Bengtsson, L., Schöneberg, T., Otto, H. and Hucho, F. 2001. Nuclear envelope proteomics: novel integral membrane proteins of the inner nuclear membrane. *Proc. Natl. Acad. Sci. USA* 98: 11943-11948.
2. Bray, J.D., Chennathukuzhi, V.M. and Hecht, N.B. 2002. Identification and characterization of cDNAs encoding four novel proteins that interact with translin associated factor-X. *Genomics* 79: 799-808.
3. Online Mendelian Inheritance in Man, OMIM™. 2005. Johns Hopkins University, Baltimore, MD. MIM Number: 607723. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
4. Padmakumar, V.C., Libotte, T., Lu, W., Zaim, H., Abraham, S., Noegel, A.A., Gotzmann, J., Foisner, R. and Karakesiosoglou, I. 2005. The inner nuclear membrane protein SUN1 mediates the anchorage of Nesprin-2 to the nuclear envelope. *J. Cell Sci.* 118: 3419-3430.
5. Wang, Q., Du, X., Cai, Z. and Greene, M.I. 2006. Characterization of the structures involved in localization of the SUN proteins to the nuclear envelope and the centrosome. *DNA Cell Biol.* 25: 554-562.
6. Hasan, S., Güttinger, S., Mühlhäusser, P., Anderegg, F., Bürgler, S. and Kutay, U. 2006. Nuclear envelope localization of human UNC84A does not require nuclear lamins. *FEBS Lett.* 580: 1263-1268.

CHROMOSOMAL LOCATION

Genetic locus: SUN1 (human) mapping to 7p22.3; Sun1 (mouse) mapping to 5 G2.

SOURCE

SUN1 (N-19) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of SUN1 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-54447 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

SUN1 (N-19) is recommended for detection of SUN1 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

SUN1 (N-19) is also recommended for detection of SUN1 in additional species, including equine, canine and porcine.

Suitable for use as control antibody for SUN1 siRNA (h): sc-106672, SUN1 siRNA (m): sc-108011, SUN1 shRNA Plasmid (h): sc-106672-SH, SUN1 shRNA Plasmid (m): sc-108011-SH, SUN1 shRNA (h) Lentiviral Particles: sc-106672-V and SUN1 shRNA (m) Lentiviral Particles: sc-108011-V.


Molecular Weight of SUN1: 100 kDa.

Positive Controls: SK-N-MC nuclear extract: sc-2154.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



SUN1 (N-19): sc-54447. Western blot analysis of SUN1 expression in SK-N-MC nuclear extract.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

MONOS
Satisfaction
Guaranteed

Try **SUN1 (2D10): sc-293292**, our highly recommended monoclonal alternative to SUN1 (N-19).