



MRP7 (P-18): sc-54545

BACKGROUND

The MRP family is represented by nine similar ABC transporters that have the ability to transport structurally diverse lipophilic anions and operate as chemical efflux pumps. MRP7 (multidrug resistance-associated protein 7, ATP-binding cassette sub-family C member 10) is a multi-pass membrane protein that belongs to the ABC transporter family (conjugate transporter subfamily). MRP7 is involved with the ATP-dependent transport of 17 β -estradiol-D-17- β -glucuronide (E₂17 β G). MRP7 is also probably involved in cellular detoxification through its lipophilic anion extrusion capabilities. MRP7 contains two ABC transmembrane type 1 domains and two ABC transporter domains. MRP7 likely has three isoforms. Isoform 2 is the most widely expressed, while isoform 1 is predominately expressed in the spleen.

REFERENCES

- Hopper, E., Belinsky, M.G., Zeng, H., Tosolini, A., Testa, J.R. and Kruh, G.D. 2001. Analysis of the structure and expression pattern of MRP7 (ABCC10), a new member of the MRP subfamily. *Cancer Lett.* 162: 181-191.
- Chen, Z.S., Hopper-Borge, E., Belinsky, M.G., Shchaveleva, I., Kotova, E. and Kruh, G.D. 2003. Characterization of the transport properties of human multidrug resistance protein 7 (MRP7, ABCC10). *Mol. Pharmacol.* 63: 351-358.
- Hopper-Borge, E., Chen, Z.S., Shchaveleva, I., Belinsky, M.G. and Kruh, G.D. 2004. Analysis of the drug resistance profile of multidrug resistance protein 7 (ABCC10): resistance to docetaxel. *Cancer Res.* 64: 4927-4930.
- Dabrowska, M. and Sirotnak, F.M. 2004. Regulation of transcription of the human MRP7 gene. Characteristics of the basal promoter and identification of tumor-derived transcripts encoding additional 5' end heterogeneity. *Gene* 341: 129-139.
- Maher, J.M., Slitt, A.L., Cherrington, N.J., Cheng, X. and Klaassen, C.D. 2005. Tissue distribution and hepatic and renal ontogeny of the multidrug resistance-associated protein (Mrp) family in mice. *Drug Metab. Dispos.* 33: 947-955.
- Weaver, D.A., Crawford, E.L., Warner, K.A., Elkhairi, F., Khuder, S.A. and Willey, J.C. 2005. ABCC5, ERCC2, XPA and XRCC1 transcript abundance levels correlate with cisplatin chemoresistance in non-small cell lung cancer cell lines. *Mol. Cancer* 4: 18-18.
- Wooden, S.L., Kalb, S.R., Cotter, R.J. and Soloski, M.J. 2005. Cutting edge: HLA-E binds a peptide derived from the ATP-binding cassette transporter multidrug resistance-associated protein 7 and inhibits NK cell-mediated lysis. *J. Immunol.* 175: 1383-1387.
- Naramoto, H., Uematsu, T., Uchihashi, T., Doto, R., Matsuura, T., Usui, Y., Uematsu, S., Li, X., Takahashi, M., Yamaoka, M. and Furusawa, K. 2007. Multidrug resistance-associated protein 7 expression is involved in cross-resistance to docetaxel in salivary gland adenocarcinoma cell lines. *Int. J. Oncol.* 30: 393-401.
- Kruh, G.D., Guo, Y., Hopper-Borge, E., Belinsky, M.G. and Chen, Z.S. 2007. ABCC10, ABCC11, and ABCC12. *Pflugers Arch.* 453: 675-684.

CHROMOSOMAL LOCATION

Genetic locus: Abcc10 (mouse) mapping to 17 B3.

SOURCE

MRP7 (P-18) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of MRP7 of mouse origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-54545 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

MRP7 (P-18) is recommended for detection of MRP7 of mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for MRP7 siRNA (m): sc-62642.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.