

MICA/B (G-18): sc-5461

BACKGROUND

MICA and MICB are stress induced antigens that are related to major histocompatibility complex (MHC) class I molecules. MICA and MICB are frequently expressed in epithelial tumors. These highly glycosylated cell surface proteins are stably expressed without conventional class I peptide ligands or association with β -2-Microglobulin. The expression is induced on proliferating or heat shock stressed epithelial cells. MICA and MICB are broadly recognized by intestinal epithelial V δ 1 $\gamma\delta$ T cells expressing variable TCRs, suggesting that these antigens may play a central role in the signaling of cellular distress to evoke immune responses in the intestinal epithelium.

REFERENCES

1. Bahram, S., et al. 1994. A second lineage of mammalian major histocompatibility complex class I genes. *Proc. Natl. Acad. Sci. USA* 91: 6259-6263.
2. Bahram, S., et al. 1996. Nucleotide sequence of the human MHC class I MICA gene. *Immunogenetics* 44: 80-81.
3. Bahram, S., et al. 1996. Genomic structure of the human MHC class I MICB gene. *Immunogenetics* 45: 161-162.
4. Groh, V., et al. 1996. Cell stress-regulated human major histocompatibility complex class I gene expressed in gastrointestinal epithelium. *Proc. Natl. Acad. Sci. USA* 93: 12445-12450.
5. Groh, V., et al. 1998. Recognition of stress-induced MHC molecules by intestinal epithelial $\gamma\delta$ T cells. *Science* 279: 1737-1740.
6. Steinle, A., et al. 1998. Diversification, expression and $\gamma\delta$ T cell recognition of evolutionarily distant members of the MIC family of major histocompatibility complex class I-related molecules. *Proc. Natl. Acad. Sci. USA* 95: 12510-12515.
7. Groh, V., et al. 1999. Broad tumor-associated expression and recognition by tumor-derived $\gamma\delta$ T cells of MICA and MICB. *Proc. Natl. Acad. Sci. USA* 96: 6879-6884.

CHROMOSOMAL LOCATION

Genetic locus: MICA/MICB (human) mapping to 6p21.33.

SOURCE

MICA/B (G-18) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of MICA of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-5461 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

APPLICATIONS

MICA/B (G-18) is recommended for detection of MICA and MICB of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for MICA/B siRNA (h): sc-43931, MICA/B shRNA Plasmid (h): sc-43931-SH and MICA/B shRNA (h) Lentiviral Particles: sc-43931-V.

Molecular Weight of truncated MICA/B: 38 kDa.


Molecular Weight of glycosylated MICA/B: 62 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200, MCF7 whole cell lysate: sc-2206 or Daudi cell lysate: sc-2415.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941. 3) Immunohistochemistry: use ImmunoCruz™: sc-2053 or ABC: sc-2023 goat IgG Staining Systems.

DATA



MICA/B (G-18): sc-5461. Immunoperoxidase staining of formalin fixed, paraffin-embedded human pancreas tissue showing cytoplasmic staining of islets of Langerhans.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.