



CA XI (C-12): sc-54744

BACKGROUND

Carbonic anhydrases (CAs) are members of a large family of zinc metalloenzymes responsible for catalyzing the reversible hydration of carbon dioxide. CAs show extensive diversity in their distribution and subcellular localization. They are involved in a variety of biological processes, including calcification, bone resorption, respiration, acid-base balance and the formation of aqueous humor, saliva, gastric juice and cerebrospinal fluid. CA XI, also referred to as carbonic anhydrase-related protein 11 precursor (CA-RP XI) or carbonic anhydrase-related protein 2 (CA-RP II), is a member of the carbonic anhydrase family that lacks two of the three Zn-binding motifs essential for carbonic anhydrase activity. For this reason, CA XI does not exhibit catalytic activity. It is expressed primarily in brain but is also found in spinal cord and thyroid. CA XI may play a role in brain development.

REFERENCES

- Smith, K.S., Jakubzick, C., Whittam, T.S. and Ferry, J.G. 2000. Carbonic anhydrase is an ancient enzyme widespread in prokaryotes. *Proc. Natl. Acad. Sci. USA* 96: 15184-15189.
- Okamoto, N., Fujikawa-Adachi, K., Nishimori, I., Taniuchi, K. and Onishi, S. 2001. cDNA sequence of human carbonic anhydrase-related protein, CA-RP X: mRNA expressions of CA-RP X and XI in human brain. *Biochim. Biophys. Acta* 1518: 311-316.
- Supuran, C.T., Scozzafava, A. and Casini, A. 2002. Carbonic anhydrase inhibitors. *Med. Res. Rev.* 23: 146-189.
- Taniuchi, K., Nishimori, I., Takeuchi, T., Ohtsuki, Y. and Onishi, S. 2003. cDNA cloning and developmental expression of murine carbonic anhydrase-related proteins VIII, X, and XI. *Brain Res. Mol. Brain Res.* 109: 207-215.
- Supuran, C.T., Vullo, D., Manole, G., Casini, A. and Scozzafava, A. 2004. Designing of novel carbonic anhydrase inhibitors and activators. *Curr. Med. Chem. Cardiovasc. Hematol. Agents* 2: 49-68.
- Pastorekova, S., Parkkila, S., Pastorek, J. and Supuran, C.T. 2004. Carbonic anhydrases: current state of the art, therapeutic applications and future prospects. *J. Enzyme Inhib. Med. Chem.* 19: 199-229.
- Morimoto, K., Nishimori, I., Takeuchi, T., Kohsaki, T., Okamoto, N., Taguchi, T., Yunoki, S., Watanabe, R., Ohtsuki, Y. and Onishi, S. 2005. Overexpression of carbonic anhydrase-related protein XI promotes proliferation and invasion of gastrointestinal stromal tumors. *Virchows Arch.* 447: 66-73.
- Supuran, C.T. and Scozzafava, A. 2007. Carbonic anhydrases as targets for medicinal chemistry. *Bioorg. Med. Chem.* 15: 4336-4350.
- Ohradanova, A., Vullo, D., Kopacek, J., Temperini, C., Betakova, T., Pastorekova, S., Pastorek, J. and Supuran, C.T. 2007. Reconstitution of carbonic anhydrase activity of the cell surface binding protein of vaccinia virus. *Biochem. J.* 407: 61-67.

CHROMOSOMAL LOCATION

Genetic locus: CA11 (human) mapping to 19q13.3; Car11 (mouse) mapping to 7 B2.

SOURCE

CA XI (C-12) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of CA XI of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-54744 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

CA XI (C-12) is recommended for detection of CA XI of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for CA XI siRNA (h): sc-62042 and CA XI siRNA (m): sc-62043.

Molecular Weight of CA XI: 36 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.