

Gas2 (Y-19): sc-54809

BACKGROUND

Gas2 is a 313 amino acid protein encoded by the human gene GAS2. Gas2 is thought to play a role in apoptosis by acting as a cell death substrate for caspases. Gas2, a component of the microfilament system, is cleaved by a caspase (caspase-3 and caspase-7) at Asparagine 278 during apoptosis. The cleaved form resulting from this dramatically induces the rearrangement of the Actin cytoskeleton and causes potent changes in the shape of the affected cells. Gas2 is believed to also be involved in the membrane ruffling process. During the G₀-G₁ transition phase Gas2 can be found phosphorylated on its serine residues. Gas2 is a cytoskeleton and peripheral membrane protein that co-localizes with Actin fibers at the cell border and along the stress fibers in growth-arrested fibroblasts. Gas2 is mainly membrane-associated but when hyperphosphorylated it will accumulate at membrane ruffles. Gas2 is specifically expressed at growth arrest and is ubiquitously expressed with highest levels found in liver, lung and kidney. There is no evidence, however, of Gas2 expression in spleen.

REFERENCES

1. Fleming, J.V., Hay, S.M., Harries, D.N. and Rees, W.D. 1998. Effects of nutrient deprivation and differentiation on the expression of growth-arrest genes (gas and gadd) in F9 embryonal carcinoma cells. *Biochem. J.* 330: 573-579
2. Collavin, L., Buzzai, M., Saccone, S., Bernard, L., Federico, C., DellaValle, G., Brancolini, C. and Schneider, C. 1998. cDNA characterization and chromosome mapping of the human GAS2 gene. *Genomics* 48: 265-269.
3. Sgorbissa, A., Benetti, R., Marzinotto, S., Schneider, C. and Brancolini, C. 2000. Caspase-3 and caspase-7 but not caspase-6 cleave Gas2 *in vitro*: implications for microfilament reorganization during apoptosis. *J. Cell Sci.* 112: 4475-4482
4. Benetti, R., Del Sal, G., Monte, M., Paroni, G., Brancolini, C. and Schneider, C. 2001. The death substrate Gas2 binds μ -calpain and increases susceptibility to p53-dependent apoptosis. *EMBO J.* 20: 2702-2714.
5. Goriounov, D., Leung, C.L. and Liem, R.K. 2003. Protein products of human Gas2-related genes on chromosomes 17 and 22 (hGAR17 and hGAR22) associate with both microfilaments and microtubules. *J. Cell Sci.* 116: 1045-1058.
6. Brockman, J.L. and Schuler, L.A. 2005. Prolactin signals via Stat5 and Oct-1 to the proximal cyclin D1 promoter. *Mol. Cell. Endocrinol.* 239: 45-53.
7. Ragni, E., Fontaine, T., Gissi, C., Latgè, J.P. and Popolo, L. 2007. The Gas family of proteins of *Saccharomyces cerevisiae*: characterization and evolutionary analysis. *Yeast* 24: 297-308.
8. Ragni, E., Coluccio, A., Rolli, E., Rodriguez-Peña, J.M., Colasante, G., Arroyo, J., Neiman, A.M. and Popolo, L. 2007. GAS2 and GAS4, a pair of developmentally regulated genes required for spore wall assembly in *Saccharomyces cerevisiae*. *Eukaryotic Cell* 6: 302-316.

CHROMOSOMAL LOCATION

Genetic locus: GAS2 (human) mapping to 11p14.3; Gas2 (mouse) mapping to 7 B5.

SOURCE

Gas2 (Y-19) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of Gas2 of mouse origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-54809 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

Gas2 (Y-19) is recommended for detection of Gas 2 of mouse and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Gas2 (Y-19) is also recommended for detection of Gas 2 in additional species, including equine, canine, bovine, porcine and avian.


Suitable for use as control antibody for Gas2 siRNA (h): sc-62368, Gas2 siRNA (m): sc-62369, Gas2 shRNA Plasmid (h): sc-62368-SH, Gas2 shRNA Plasmid (m): sc-62369-SH, Gas2 shRNA (h) Lentiviral Particles: sc-62368-V and Gas2 shRNA (m) Lentiviral Particles: sc-62369-V.

Molecular Weight (predicted) of Gas2: 35 kDa.

Molecular Weight (observed) of Gas2: 30-37 kDa.

Positive Controls: NIH/3T3 whole cell lysate: sc-2210, Hep G2 cell lysate: sc-2227 or A549 cell lysate: sc-2413.

DATA



Gas2 (Y-19): sc-54809. Western blot analysis of Gas2 expression in HeLa (A), A549 (B), CCD-1064Sk (C), NIH/3T3 (D) and Hep G2 (E) whole cell lysates.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.