

GMAP-210 (C-16): sc-54812

BACKGROUND

Golgi microtubule-associated protein-210 (GMAP-210), also referred to as CEV14, Trip11 or Trip230, is a peripheral Golgi protein that localizes to the *cis*-Golgi network. GMAP-210 is a 1,978 amino acid coiled-coil member of the golgin family of proteins. Microtubule ends bind to GMAP-210 which functions to link the *cis*-Golgi network to the minus ends of centrosome-nucleated microtubules. This interaction may be essential for the proper morphology and structural maintenance of the Golgi apparatus. GMAP-210 also associates with thyroid hormone receptor β . Overexpression of GMAP-210 disrupts the microtubule network and causes a significant enlargement and fragmentation of the Golgi apparatus; it also blocks anterograde and retrograde transport between the ER and the Golgi apparatus.

REFERENCES

- Infante, C., Ramos-Morales, F., Fedriani, C., Bornens, M. and Rios, R.M. 1999. GMAP-210, *cis*-Golgi network-associated protein, is a minus end microtubule-binding protein. *J. Cell Biol.* 145: 83-98.
- Ramos-Morales, F., Vime, C., Bornens, M., Fedriani, C. and Rios, R.M. 2001. Two splice variants of Golgi microtubule-associated protein of 210 kDa (GMAP-210) differ in their binding to the *cis*-Golgi network. *Biochem. J.* 357: 699-708.
- Pernet-Gallay, K., Antony, C., Johannes, L., Bornens, M., Goud, B. and Rios, R.M. 2002. The overexpression of GMAP-210 blocks anterograde and retrograde transport between the ER and the Golgi apparatus. *Traffic* 3: 822-832.
- Online Mendelian Inheritance in Man, OMIM[™]. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 604505. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- Gillingham, A.K., Tong, A.H., Boone, C. and Munro, S. 2004. The GTPase Arf1p and the ER to Golgi cargo receptor Erv14p cooperate to recruit the golgin Rud3p to the *cis*-Golgi. *J. Cell Biol.* 167: 281-292.
- Linstedt, A.D. 2004. Positioning the Golgi apparatus. *Cell* 118: 271-272.
- Ríos, R.M., Sanchís, A., Tassin, A.M., Fedriani, C. and Bornens, M. 2004. GMAP-210 recruits γ Tubulin complexes to *cis*-Golgi membranes and is required for Golgi ribbon formation. *Cell* 118: 323-335.
- Barr, F.A. and Egerer, J. 2005. Golgi positioning: are we looking at the right MAP? *J. Cell Biol.* 168: 993-998.
- Ouchi, M., West, K., Crabb, J.W., Kinoshita, S. and Kamei, M. 2005. Proteomic analysis of vitreous from diabetic macular edema. *Exp. Eye Res.* 81: 176-182.

CHROMOSOMAL LOCATION

Genetic locus: TRIP11 (human) mapping to 14q32.12; Trip11 (mouse) mapping to 12 F1.

SOURCE

GMAP-210 (C-16) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of GMAP-210 of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-54812 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

GMAP-210 (C-16) is recommended for detection of GMAP-210 of human origin and, to a lesser extent, TRIP11 of mouse and rat origin of human by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for GMAP-210 siRNA (h): sc-62387, TRIP11 siRNA (m): sc-154677, GMAP-210 shRNA Plasmid (h): sc-62387-SH, TRIP11 shRNA Plasmid (m): sc-154677-SH, GMAP-210 shRNA (h) Lentiviral Particles: sc-62387-V and TRIP11 shRNA (m) Lentiviral Particles: sc-154677-V.

Molecular Weight of GMAP-210: 210 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker[™] compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker[™] Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz[™] Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.


 MONOS
 Satisfaction
 Guaranteed

Try **GMAP-210 (E-2): sc-515208** or **GMAP-210 (15): sc-135928**, our highly recommended monoclonal alternatives to GMAP-210 (C-16).