

ABC1 (N-15): sc-5491

BACKGROUND

ABC1 (for ATP-binding cassette transporter 1) is a member of the family of ATP-binding cassette proteins which transport various molecules across biological membranes. ABC1 contains 2 characteristic ATP-binding domains and 12 transmembrane domains which form a channel-like structure for transport. Mutations in the ABC1 gene are implicated in Tangier disease, characterized by low serum high density lipoprotein. ABC1 is widely expressed in human tissues, with high levels of expression in liver, lung, adrenal glands, placenta and fetal tissue. ABC1 expression is induced during monocyte differentiation and upregulated in the presence of acetylated low-density lipoprotein. ABC1 may have a dual regulatory function in macrophage lipid metabolism and inflammation.

REFERENCES

1. Decottignies, A. and Goffeau, A. 1997. Complete inventory of the yeast ABC proteins. *Nat. Genet.* 15: 137-145.
2. Schwiebert, E.M. 1999. ABC transporter-facilitated ATP conductive transport. *Am. J. Physiol.* 276: C1-C8.

CHROMOSOMAL LOCATION

Genetic locus: ABCA1 (human) mapping to 9q31.1; Abca1 (mouse) mapping to 4 B2.

SOURCE

ABC1 (N-15) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of ABC1 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-5491 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

ABC1 (N-15) is recommended for detection of ABC1 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

ABC1 (N-15) is also recommended for detection of ABC1 in additional species, including equine, canine, bovine, porcine and avian.

Suitable for use as control antibody for ABC1 siRNA (h): sc-41136, ABC1 siRNA (m): sc-41137, ABC1 shRNA Plasmid (h): sc-41136-SH, ABC1 shRNA Plasmid (m): sc-41137-SH, ABC1 shRNA (h) Lentiviral Particles: sc-41136-V and ABC1 shRNA (m) Lentiviral Particles: sc-41137-V.


Molecular Weight of ABC1: 220 kDa.

Positive Controls: MES-SA/Dx5 cell lysate: sc-2284.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



ABC1 (N-15): sc-5491. Western blot analysis of ABC1 expression in MES-SA/Dx5 whole cell lysate.



ABC1 (N-15): sc-5491. Immunofluorescence staining of methanol-fixed RAW 264.7 cells showing cytoplasmic localization.

SELECT PRODUCT CITATIONS

1. Chang, H., et al. 2004. Using siRNA technique to generate transgenic animals with spatiotemporal and conditional gene knockdown. *Am. J. Pathol.* 165: 1535-1541.
2. Baker, A.D., et al. 2010. Targeted PPAR γ deficiency in alveolar macrophages disrupts surfactant catabolism. *J. Lipid Res.* 51: 1325-1331.
3. Collaco, R.C., et al. 2012. Protection by *Mikania laevigata* (guaco) extract against the toxicity of *Philodryas olfersii* snake venom. *Toxicon* 60: 614-622.
4. Mostafa, A.M., et al. 2015. Glucagon-like peptide 1 (GLP-1)-based therapy upregulates LXR-ABCA1/ABCG1 cascade in adipocytes. *Biochem. Biophys. Res. Commun.* E-Published.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

MONOS
Satisfaction
Guaranteed

Try **ABC1 (AB.H10): sc-58219**, our highly recommended monoclonal alternative to ABC1 (N-15).