

PIG-S (S-15): sc-54982

BACKGROUND

Phosphatidylinositol-glycans (PIGs) are multi-pass transmembrane proteins that localize to the endoplasmic reticulum. PIGs exhibit various functions but all are crucial for the biosynthesis of the glycosylphosphatidylinositol (GPI)-anchor. Some PIG proteins are components of the GPI transamidase (GPIT) complex and play a role in the recognition of either the GPI attachment signal or the lipid portion of GPI. Other PIGs belong to the glycosyltransferase complex and function in the transfer of N-acetylglucosamine (GlcNAc) to phosphatidylinositol (PI). A variety of other PIGs play distinct roles in GPI synthesis. PIG-S is a component of GPIT, a multisubunit membrane-bound complex that recognizes the C-terminal signal sequences on proproteins, cleaves them and replaces them with specific GPI lipids. PIG-S is required for the generation of the carbonyl intermediate.

REFERENCES

- Ohishi, K., Inoue, N. and Kinoshita, T. 2001. PIG-S and PIG-T, essential for GPI anchor attachment to proteins, form a complex with GAA1 and GPI8. *EMBO J.* 20: 4088-4098.
- Hong, Y., Ohishi, K., Kang, J.Y., Tanaka, S., Inoue, N., Nishimura, J., Maeda, Y. and Kinoshita, T. 2003. Human PIG-U and yeast Cdc91p are the fifth subunit of GPI transamidase that attaches GPI-anchors to proteins. *Mol. Biol. Cell* 14: 1780-1789.
- Nagamune, K., Ohishi, K., Ashida, H., Hong, Y., Hino, J., Kangawa, K., Inoue, N., Maeda, Y. and Kinoshita, T. 2003. GPI transamidase of *Trypanosoma brucei* has two previously uncharacterized (trypanosomatid transamidase 1 and 2) and three common subunits. *Proc. Natl. Acad. Sci. USA* 100: 10682-10687.
- Vainauskas, S. and Menon, A.K. 2004. A conserved proline in the last transmembrane segment of Gaa1 is required for glycosylphosphatidylinositol (GPI) recognition by GPI transamidase. *J. Biol. Chem.* 279: 6540-6545.
- Zhu, Y., Fraering, P., Vionnet, C. and Conzelmann, A. 2005. Gpi17p does not stably interact with other subunits of glycosylphosphatidylinositol transamidase in *Saccharomyces cerevisiae*. *Biochim. Biophys. Acta* 1735: 79-88.
- Vainauskas, S. and Menon, A.K. 2006. Ethanolamine phosphate linked to the first mannose residue of glycosylphosphatidylinositol (GPI) lipids is a major feature of the GPI structure that is recognized by human GPI transamidase. *J. Biol. Chem.* 281: 38358-38364.
- Bowman, S.M., Piwowar, A., Al Dabbous, M., Vierula, J. and Free, S.J. 2006. Mutational analysis of the glycosylphosphatidylinositol (GPI) anchor pathway demonstrates that GPI-anchored proteins are required for cell wall biogenesis and normal hyphal growth in *Neurospora crassa*. *Eukaryotic Cell* 5: 587-600.
- Hong, Y., Nagamune, K., Ohishi, K., Morita, Y.S., Ashida, H., Maeda, Y. and Kinoshita, T. 2006. TbGPI16 is an essential component of GPI transamidase in *Trypanosoma brucei*. *FEBS Lett.* 580: 603-606.
- Pittet, M. and Conzelmann, A. 2007. Biosynthesis and function of GPI proteins in the yeast *Saccharomyces cerevisiae*. *Biochim. Biophys. Acta* 1771: 405-420.

CHROMOSOMAL LOCATION

Genetic locus: PIGS (human) mapping to 17p11.2; Pigs (mouse) mapping to 11 B5.

SOURCE

PIG-S (S-15) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of PIG-S of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-54982 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

PIG-S (S-15) is recommended for detection of PIG-S of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

PIG-S (S-15) is also recommended for detection of PIG-S in additional species, including equine and porcine.

Suitable for use as control antibody for PIG-S siRNA (h): sc-62808, PIG-S siRNA (m): sc-62809, PIG-S shRNA Plasmid (h): sc-62808-SH, PIG-S shRNA Plasmid (m): sc-62809-SH, PIG-S shRNA (h) Lentiviral Particles: sc-62808-V and PIG-S shRNA (m) Lentiviral Particles: sc-62809-V.

Molecular Weight of PIG-S: 65 kDa.

Positive Controls: A-431 whole cell lysate: sc-2201 or NIH/3T3 whole cell lysate: sc-2210.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.