

NFAT5 (C-19): sc-5499

BACKGROUND

Members of the NFAT (nuclear factor of activated T cells) family of transcription factors are related to NF κ B/Rel proteins and form cooperative complexes with the AP-1 proteins, Fos and Jun, on DNA to regulate cytokine expression in T cells. NFAT proteins are widely expressed and alternatively modified to generate splice variants, and they are localized to both the cytosol (NFATc) and to the nucleus (NFATn). NFAT1, NFAT2, and NFAT4 are predominantly expressed in immune cells, and NFAT2 and NFAT3 are expressed at high levels in cardiac tissues. In addition to activating cytokine gene transcription, NFAT2 is also implicated in cardiac valve development, and NFAT3 is involved in cardiac hypertrophy. NFAT5 is detected in both immune and nonimmune cells and, like other NFAT proteins, contains a highly conserved Rel-like binding domain that mediates NFAT proteins associating with specific consensus sequences on DNA. NFAT proteins are activated by increases in intracellular calcium, which leads to the calmodulin-dependent phosphatase, calcineurin, dephosphorylating NFAT proteins. This activating event induces a conformational change in the protein structure that exposes the nuclear localization signal and facilitates the translocation of NFAT proteins from the cytosol into the nucleus.

REFERENCES

1. Hoey, T., et al. 1995. Isolation of two new members of the NFAT gene family and functional characterization of the NFAT proteins. *Immunity* 2: 461-472.
2. Park, J., et al. 1996. Characterization of a new isoform of the NFAT (nuclear factor of activated T cells) gene family member NFATc. *J. Biol. Chem.* 271: 20914-20921.
3. Rao, A., et al. 1997. Transcription factors of the NFAT family: regulation and function. *Annu. Rev. Immunol.* 15: 707-747.
4. Lyakh, L., et al. 1997. Expression of NFAT-family proteins in normal human T cells. *Mol. Cell. Biol.* 17: 2475-2484.
5. Ranger, A.M., et al. 1998. The transcription factor NFATc is essential for cardiac valve formation. *Nature* 392: 186-190.

CHROMOSOMAL LOCATION

Genetic locus: NFAT5 (human) mapping to 16q22.1; Nfat5 (mouse) mapping to 8 D2.

SOURCE

NFAT5 (C-19) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the C-terminus of NFAT5 of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin. Also available as TransCruz reagent for Gel Supershift and ChIP applications, sc-5499 X, 200 μ g/0.1 ml.

Blocking peptide available for competition studies, sc-5499 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

NFAT5 (C-19) is recommended for detection of NFAT5a, NFAT5b and NFAT5c of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000)

NFAT5 (C-19) is also recommended for detection of NFAT5a, NFAT5b and NFAT5c in additional species, including equine, canine, bovine, porcine and avian.

Suitable for use as control antibody for NFAT5 siRNA (h): sc-43968, NFAT5 siRNA (m): sc-38122, NFAT5 shRNA Plasmid (h): sc-43968-SH, NFAT5 shRNA Plasmid (m): sc-38122-SH, NFAT5 shRNA (h) Lentiviral Particles: sc-43968-V and NFAT5 shRNA (m) Lentiviral Particles: sc-38122-V.

NFAT5 (C-19) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.

Molecular Weight of NFAT5: 170 kDa.

Positive Controls: Jurkat whole cell lysate: sc-2204, HeLa whole cell lysate: sc-2200 or IMR-32 cell lysate: sc-2409.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



Try **NFAT5 (F-9): sc-398171** or **NFAT5 (37X): sc-101098**, our highly recommended monoclonal alternatives to NFAT5 (C-19).