# Aminoacylase-1 (L-19): sc-55060



The Power to Question

## **BACKGROUND**

Aminoacylase-1, also designated N-acyl-L-amino-acid amidohydrolase or ACY-1, is a member of the largest metallopeptidase family called M20A. Aminoacylase-1 is a zinc-binding homodimeric enzyme expressed in kidney, brain, placenta and spleen. It is the most abundant of the aminoacylases. Aminoacylase-1 localizes to the cytoplasm and catalyzes the hydrolysis of N-acylated or N-acetylated amino acids. In addition, Aminoacylase-1 is a sphingosine kinase 1 (SphK1)-interacting protein and may also play a role in regulating responses of the cell to oxidative stress. The gene encoding Aminoacylase-1 is evolutionarily conserved in fish, frog, mouse, rat and human. Deficiency in the Aminoacylase-1 protein may result in defects of brain metabolism and function.

## **REFERENCES**

- Cook, R.M., et al. 1993. Human Aminoacylase-1. Cloning, sequence, and expression analysis of a chromosome 3p21 gene inactivated in small cell lung cancer. J. Biol. Chem. 268: 17010-17017.
- Lindner, H., et al. 2001. Mutational analysis of two PWW sequence motifs in human aminoacylase 1. Biol. Chem. 381: 1055-1061.
- Lindner, H.A., et al. 2003. Essential roles of zinc ligation and enzyme dimerization for catalysis in the aminoacylase-1/M20 family. J. Biol. Chem. 278: 44496-44504.
- Maceyka, M., et al. 2004. Aminoacylase-1 is a sphingosine kinase 1-interacting protein. FEBS Lett. 568: 30-34.
- 5. Lindner, H.A., et al. 2005. Roles of dimerization domain residues in binding and catalysis by Aminoacylase-1. Biochemistry 44: 15645-15651.
- Liu, Z., et al. 2006. Probing the catalytic center of porcine Aminoacylase-1 by site-directed mutagenesis, homology modeling and substrate docking. J. Biochem. 139: 421-430.
- Sass, J.O., et al. 2006. Mutations in ACY, the gene encoding Aminoacylase-1, cause a novel inborn error of metabolism. Am. J. Hum. Genet. 78: 401-409.
- Engelke, U.F., et al. 2007. NMR spectroscopy of Aminoacylase-1 deficiency, a novel inborn error of metabolism. NMR Biomed. 21: 138-147.

# CHROMOSOMAL LOCATION

Genetic locus: ACY1 (human) mapping to 3p21.2.

## SOURCE

Aminoacylase-1 (L-19) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of Aminoacylase-1 of human origin.

## **PRODUCT**

Each vial contains 200  $\mu g$  lgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-55060 P, (100  $\mu$ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

#### **APPLICATIONS**

Aminoacylase-1 (L-19) is recommended for detection of Aminoacylase-1 of human and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2  $\mu$ g per 100-500  $\mu$ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Aminoacylase-1 (L-19) is also recommended for detection of Aminoacylase-1 in additional species, including equine, canine and bovine.


Suitable for use as control antibody for Aminoacylase-1 siRNA (h): sc-61966, Aminoacylase-1 shRNA Plasmid (h): sc-61966-SH and Aminoacylase-1 shRNA (h) Lentiviral Particles: sc-61966-V.


Molecular Weight (predicted) of Aminoacylase-1: 45 kDa.


Molecular Weight (observed) of Aminoacylase-1: 42 kDa.

Positive Controls: K-562 whole cell lysate: sc-2203, Caki-1 cell lysate: sc-2224 or HeLa whole cell lysate: sc-2200.

#### **DATA**







Aminoacylase-1 (L-19): sc-55060. Immunofluorescence staining of methanol-fixed HeLa cells showing cytoplasmic localization.

## **STORAGE**

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## **RESEARCH USE**

For research use only, not for use in diagnostic procedures.

## **PROTOCOLS**

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

Santa Cruz Biotechnology, Inc. 1.800.457.3801 831.457.3801 Fax 831.457.3801 Europe +00800 4573 8000 49 6221 4503 0 www.scbt.com