

# APPL2 (N-16): sc-55069

## BACKGROUND

The APPL family of proteins are involved in linking, trafficking and signaling downstream of tyrosine kinase receptors. APPL1, also designated adaptor protein containing pH domain, PTB domain and leucine zipper motif 1; APPL or DCC interacting protein 13 $\alpha$  (DIP13 $\alpha$ ), and APPL2, also designated adaptor protein containing pH domain, PTB domain and leucine zipper motif 2 or DCC interacting protein 13 $\beta$  (DIP13 $\beta$ ), are involved in the coupling of epidermal growth factor (EGF) signaling and chromatin remodeling in the nucleus. They associate with GTPase Rab 5 and are released from the plasma membrane and translocated to the nucleus. In the nucleus, APPL1 and APPL2 associate with NuRD/MeCP1 and are essential for cell growth and proliferation. APPL2 also associates with follicle stimulating hormone receptor (FSHR). APPL2 is highly expressed in heart, brain, skeletal muscle, and kidney. APPL2 shares 54% homology with APPL1.

## CHROMOSOMAL LOCATION

Genetic locus: APPL2 (human) mapping to 12q23.3; Appl2 (mouse) mapping to 10 C1.

## SOURCE

APPL2 (N-16) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the N-terminus of APPL2 of human origin.

## PRODUCT

Each vial contains 200  $\mu$ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-55069 P, (100  $\mu$ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## STORAGE

Store at 4 $^{\circ}$  C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## APPLICATIONS

APPL2 (N-16) is recommended for detection of APPL2 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2  $\mu$ g per 100-500  $\mu$ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

APPL2 (N-16) is also recommended for detection of APPL2 in additional species, including equine, canine, bovine, porcine and avian.

Suitable for use as control antibody for APPL2 siRNA (h): sc-61982, APPL2 siRNA (m): sc-61983, APPL2 shRNA Plasmid (h): sc-61982-SH, APPL2 shRNA Plasmid (m): sc-61983-SH, APPL2 shRNA (h) Lentiviral Particles: sc-61982-V and APPL2 shRNA (m) Lentiviral Particles: sc-61983-V.


Molecular Weight of APPL2: 80 kDa.

Positive Controls: APPL2 (m): 293T Lysate: sc-118502 or Hs 181 Tes whole cell lysate: sc-364779.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## DATA



APPL2 (N-16): sc-55069. Western blot analysis of APPL2 expression in non-transfected: sc-117752 (A) and mouse APPL2 transfected: sc-118502 (B) 293T whole cell lysates.

## SELECT PRODUCT CITATIONS

1. Tan, Y., et al. 2009. Appl1 is dispensable for mouse development, and loss of Appl1 has growth factor-selective effects on Akt signaling in murine embryonic fibroblasts. *J. Biol. Chem.* 285: 6377-6389.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.

**MONOS**  
Satisfaction  
Guaranteed

Try **APPL2 (F-2): sc-398859** or **APPL2 (E-1): sc-271084**, our highly recommended monoclonal alternatives to APPL2 (N-16).