

DGCR6 (N-16): sc-55091

BACKGROUND

Neural crest cell migration to the third and fourth pharyngeal pouches is a critical step in the structural formation of organs that are affected in DiGeorge syndrome, a developmental disorder characterized by improper facial, cardiac and palate formation. DGCR6 (DiGeorge syndrome critical region 6) is a nuclear protein that plays a role in neural crest cell migration and is located at the DiGeorge syndrome critical region (DGCR) on chromosome 22. Expressed ubiquitously with highest levels in heart, liver and skeletal muscle, DGCR6 shares high homology with the *Drosophila* gonadal (gdl) protein and with human Laminin γ -1, both of which are involved in early tissue development. The gene encoding DGCR6, along with other DGCR genes, is deleted in DiGeorge syndrome. Upregulation of DGCR6 is implicated in lung and colon adenocarcinomas, as well as in Burkitt's lymphoma and lymphocytes transformed by EBV. Due to a duplication of the ancestral DGCR6 locus, there are two functional, highly homologous copies of the DGCR6 gene (designated DGCR6 and DGCR6L) on chromosome 22.

REFERENCES

- Lindsay, E.A. and Baldini, A. 1998. A mouse gene (Dgcr6) related to the *Drosophila* gonadal gene is expressed in early embryogenesis and is the homolog of a human gene deleted in DiGeorge syndrome. *Cytogenet. Cell Genet.* 79: 243-247.
- Edelmann, L., Stankiewicz, P., Spiteri, E., Pandita, R.K., Shaffer, L., Lupski, J.R., Morrow, B.E. and Lupski, J. 2001. Two functional copies of the DGCR6 gene are present on human chromosome 22q11 due to a duplication of an ancestral locus. *Genome Res.* 11: 208-217.
- Liu, H., Heath, S.C., Sobin, C., Roos, J.L., Galke, B.L., Blundell, M.L., Lenane, M., Robertson, B., Wijsman, E.M., Rapoport, J.L., Gogos, J.A. and Karayiorgou, M. 2002. Genetic variation at the 22q11 PRODH2/DGCR6 locus presents an unusual pattern and increases susceptibility to schizophrenia. *Proc. Natl. Acad. Sci. USA* 99: 3717-3722.
- Chakravarti, A. 2002. A compelling genetic hypothesis for a complex disease: PRODH2/DGCR6 variation leads to schizophrenia susceptibility. *Proc. Natl. Acad. Sci. USA* 99: 4755-4756.
- Hierck, B.P., Molin, D.G., Boot, M.J., Poelmann, R.E. and Gittenberger-de Groot, A.C. 2004. A chicken model for DGCR6 as a modifier gene in the DiGeorge critical region. *Pediatr. Res.* 56: 440-448.
- Pfuhl, T., Dürr, M., Spurk, A., Schwalbert, B., Nord, R., Mysliwicz, J., Kremmer, E. and Grässer, F.A. 2005. Biochemical characterisation of the proteins encoded by the DiGeorge critical region 6 (DGCR6) genes. *Hum. Genet.* 117: 70-80.

CHROMOSOMAL LOCATION

Genetic locus: DGCR6/DGCR6L (human) mapping to 22q11.21.

SOURCE

DGCR6 (N-16) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of DGCR6 of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-55091 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

DGCR6 (N-16) is recommended for detection of DGCR6 and DGCR6L of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

DGCR6 (N-16) is also recommended for detection of DGCR6 and DGCR6L in additional species, including bovine.

Suitable for use as control antibody for DGCR6 siRNA (h): sc-62208, DGCR6 shRNA Plasmid (h): sc-62208-SH and DGCR6 shRNA (h) Lentiviral Particles: sc-62208-V.

Molecular Weight of DGCR6: 25 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.