

granzyme A (E-15): sc-5511

BACKGROUND

Granzyme A and granzyme B are serine proteases that mediate apoptotic signaling in cytotoxic T lymphocytes (CTL) and natural killer (NK) cells. Both granzyme A and granzyme B are synthesized as inactive proenzymes, and they are stored within cytolytic granules and released by effector cells during degranulation. In activated CTLs, granzyme A and granzyme B are processed and activated by cathepsin C, and they then function to induce apoptosis by two distinct pathways. Granzyme B proteolytically cleaves and activates members of the caspase family of cysteine proteases, including caspase-3, caspase-6, caspase-7 and caspase-9. When cleaved, these caspases assemble into active holoenzymes that then mediate apoptosis through a defined proteolytic cascade involving nuclear lamins and PARP (poly ADP ribose polymerase). Granzyme A mediates the activation of apoptosis by inducing single-strand DNA breaks, membrane perturbation and nuclear condensations in an alternative pathway that is independent from caspase activation or the caspase proteolytic cascade.

REFERENCES

1. Gershenfeld, H.K., et al. 1988. Cloning and chromosomal assignment of a human cDNA encoding a T cell- and natural killer cell-specific trypsin-like serine protease. *Proc. Natl. Acad. Sci. USA* 85: 1184-1188.
2. Shresta, S., et al. 1995. Natural killer and lymphokine-activated killer cells require granzyme B for the rapid induction of apoptosis in susceptible target cells. *Proc. Natl. Acad. Sci. USA* 92: 5679-5683.
3. Trapani, J.A., et al. 1996. A putative role in the mechanism of cytotoxic lymphocyte-mediated apoptosis. Localization of granzyme B in the nucleus. *J. Biol. Chem.* 271: 4127-4133.
4. Trapani, J.A., et al. 1998. Efficient nuclear targeting of granzyme B and the nuclear consequences of apoptosis induced by granzyme B and perforin are caspase-dependent, but cell death is caspase-independent. *J. Biol. Chem.* 273: 27934-27938.
5. Atkinson, E.A., et al. 1998. Cytotoxic T lymphocyte-assisted suicide. caspase-3 activation is primarily the result of the direct action of granzyme B. *J. Biol. Chem.* 273: 21261-21266.
6. Pham, C.T., et al. 1999. Dipeptidyl peptidase I is required for the processing and activation of granzymes A and B *in vivo*. *Proc. Natl. Acad. Sci. USA* 96: 8627-8632.
7. Shresta, S., et al. 1999. Granzyme A initiates an alternative pathway for granule-mediated apoptosis. *Immunity* 10: 595-605.

CHROMOSOMAL LOCATION

Genetic locus: GZMA (human) mapping to 5q11.2.

SOURCE

granzyme A (E-15) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of granzyme A of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-5511 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

granzyme A (E-15) is recommended for detection of granzyme A of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for granzyme A siRNA (h): sc-37431, granzyme A shRNA Plasmid (h): sc-37431-SH and granzyme A shRNA (h) Lentiviral Particles: sc-37431-V.

Molecular Weight of granzyme A monomer: 28 kDa.

Molecular Weight of granzyme A homodimer: 43-65 kDa.

Positive Controls: HL-60 whole cell lysate: sc-2209, U-937 cell lysate: sc-2239 or K-562 whole cell lysate: sc-2203.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.


 MONOS
 Satisfation
 Guaranteed

Try **granzyme A (GA6): sc-56116**, our highly recommended monoclonal alternative to granzyme A (E-15).