

# DPP7 (N-20): sc-55173

## BACKGROUND

Dipeptidyl peptidases (DPPs) mediate regulatory activity of their substrates and have been linked to a variety of diseases including type 2 diabetes, obesity and cancer. DPPs have post-proline dipeptidyl aminopeptidase activity, cleaving Xaa-Pro dipeptides from the N-termini of proteins. DPPs can bind specific voltage-gated potassium channels and alter their expression and biophysical properties and may also influence T cells. DPP proteins include DPP1, DPP2, DPP3, DPP7, DPP10, DPPX and CD26. DPP7 (dipeptidyl-peptidase 7), also known as DPP2, DPPII or QPP (quiescent cell proline dipeptidase), is expressed in quiescent lymphocytes and localizes to lysosomes. In response to calcium release, DPP7 can be secreted in its active form. DPP7 exists as a homodimer via its leucine zipper motif and is involved in the degradation of oligopeptides. DPP7 is essential for lymphocyte survival, as the inhibition of DPP7 results in quiescent cell apoptosis.

## REFERENCES

- Chiravuri, M., et al. 2000. Vesicular localization and characterization of a novel post-proline-cleaving aminodipeptidase, quiescent cell proline dipeptidase. *J. Immunol.* 165: 5695-5702.
- Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 610537. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- Rosenblum, J.S. and Kozarich, J.W. 2003. Prolyl peptidases: a serine protease subfamily with high potential for drug discovery. *Curr. Opin. Chem. Biol.* 7: 496-504. PMID: 12941425
- Hu, Y., et al. 2005. Synthesis and structure-activity relationship of N-alkyl Gly-boro-Pro inhibitors of DPP4, FAP, and DPP7. *Bioorg. Med. Chem. Lett.* 15: 4239-4242.
- Shreder, K.R., et al. 2005. Boro-norleucine as a P1 residue for the design of selective and potent DPP7 inhibitors. *Bioorg. Med. Chem. Lett.* 15: 4256-4260.
- Maes, M.B., et al. 2005. Kinetic investigation of human dipeptidyl peptidase II (DPPII)-mediated hydrolysis of dipeptide derivatives and its identification as quiescent cell proline dipeptidase (QPP)/dipeptidyl peptidase 7 (DPP7). *Biochem. J.* 386: 315-324.
- Danilova, O., et al. 2007. Synthesis and activity of a potent, specific azabicyclo[3.3.0]-octane-based DPP II inhibitor. *Bioorg. Med. Chem. Lett.* 17: 507-510.

## CHROMOSOMAL LOCATION

Genetic locus: DPP7 (human) mapping to 3p13.

## SOURCE

DPP7 (N-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the N-terminus of DPP7 of human origin.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-55173 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## APPLICATIONS

DPP7 (N-20) is recommended for detection of DPP7 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for DPP7 siRNA (h): sc-62232, DPP7 shRNA Plasmid (h): sc-62232-SH and DPP7 shRNA (h) Lentiviral Particles: sc-62232-V.

Molecular Weight of DPP7 (predicted): 54 kDa.


Molecular Weight of DPP7 (observed): 50 kDa.

Positive Controls: DPP7 (h): 293 Lysate: sc-112766.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## DATA



DPP7 (N-20): sc-55173. Western blot analysis of DPP7 expression in non-transfected 293: sc-110760 (A), HeLa (B) and human DPP7 transfected 293: sc-112766 (C) whole cell lysates.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.