

GPRC5D (S-15): sc-55371

BACKGROUND

GPRC5D (G protein-coupled receptor family C group 5 member D) is a 344 amino acid protein encoded by the mouse GPRC5D gene. GPRC5D is an orphan receptor member of the G protein-coupled receptor 3 family and a member of RAIG family. G protein-coupled receptors (GPCRs or GPRs) contain seven transmembrane domains and transduce extracellular signals through heterotrimeric G proteins. Key roles for G protein-coupled receptors include control of protein maturation and cell surface delivery and providing the correct framework for interactions with both heterotrimeric G proteins and arrestins to allow signal generation and its termination. This retinoic acid-inducible G protein-coupled receptor provides evidence for a possible interaction between retinoid and G protein signaling pathways. GPRC5D is found in hard keratinized structures.

REFERENCES

1. Bräuner-Osborne, H. and Krogsgaard-Larsen, P. 2000. Sequence and expression pattern of a novel human orphan G protein-coupled receptor, GPRC5B, a family C receptor with a short amino-terminal domain. *Genomics* 65: 121-128.
2. Robbins, M.J., Michalovich, D., Hill, J., Calver, A.R., Medhurst, A.D., Gloger, I., Sims, M., Middlemiss, D.N. and Pangalos, M.N. 2000. Molecular cloning and characterization of two novel retinoic acid-inducible orphan G protein-coupled receptors (GPRC5B and GPRC5C). *Genomics* 67: 8-18.
3. Robbins, M.J., Charles, K.J., Harrison, D.C. and Pangalos, M.N. 2002. Localisation of the GPRC5B receptor in the rat brain and spinal cord. *Brain Res. Mol. Brain Res.* 106: 136-144.
4. Takeda, S., Kadowaki, S., Haga, T., Takaesu, H. and Mitaku, S. 2002. Identification of G protein-coupled receptor genes from the human genome sequence. *FEBS Lett.* 520: 97-101.

CHROMOSOMAL LOCATION

Genetic locus: *Gprc5d* (mouse) mapping to 6 G1.

SOURCE

GPRC5D (S-15) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an N-terminal extracellular domain of GPRC5D of mouse origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-55371 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

GPRC5D (S-15) is recommended for detection of GPRC5D of mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for GPRC5D siRNA (m): sc-62412, GPRC5D shRNA Plasmid (m): sc-62412-SH and GPRC5D shRNA (m) Lentiviral Particles: sc-62412-V.

Molecular Weight (predicted) of GPRC5D isoforms: 39/34/37 kDa.


Molecular Weight (observed) of GPRC5D: 29-45 kDa.

Positive Controls: mouse kidney extract: sc-2255 or rat pancreas extract: sc-364806.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



GPRC5D (S-15): sc-55371. Western blot analysis of GPRC5D expression in rat pancreas (A) and mouse kidney (B) tissue extracts.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

MONOS
Satisfaction
Guaranteed

Try **GPRC5D (E-7): sc-377504**, our highly recommended monoclonal alternative to GPRC5D (S-15).