

caveolin-3 (C-2): sc-55518

BACKGROUND

Caveolae (also known as plasmalemmal vesicles) are 50-100 nm flask-shaped membranes that represent a subcompartment of the plasma membrane. On the basis of morphological studies, caveolae have been implicated to function in the transcytosis of various macromolecules (including LDL) across capillary endothelial cells, uptake of small molecules via potocytosis and the compartmentalization of certain signaling molecules including G protein-coupled receptors. Three proteins, caveolin-1, caveolin-2 and caveolin-3, have been identified as principal components of caveolae. Two forms of caveolin-1, designated α and β , share a distinct but overlapping cellular distribution and differ by an amino terminal 31 amino acid sequence, which is absent from the β -isoform. Caveolin-1 shares 31% identity with caveolin-2 and 65% identity with caveolin-3 at the amino acid level. Functionally, the three proteins differ in their interactions with heterotrimeric G protein isoforms.

REFERENCES

1. Fan, J.Y., et al. 1983. Morphological changes of the 3T3-L1 fibroblast plasma membrane upon differentiation to the adipocyte form. *J. Cell Sci.* 61: 219-230.
2. Rothberg, K.G., et al. 1992. Caveolin, a protein component of caveolae membrane coats. *Cell* 68: 673-682.

CHROMOSOMAL LOCATION

Genetic locus: CAV3 (human) mapping to 3p25.3; Cav3 (mouse) mapping to 6 E3.

SOURCE

caveolin-3 (C-2) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 3-40 at the N-terminus of caveolin-3 of mouse origin.

PRODUCT

Each vial contains 200 μ g IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

caveolin-3 (C-2) is available conjugated to agarose (sc-55518 AC), 500 μ g/0.25 ml agarose in 1 ml, for IP; to HRP (sc-55518 HRP), 200 μ g/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-55518 PE), fluorescein (sc-55518 FITC), Alexa Fluor[®] 488 (sc-55518 AF488), Alexa Fluor[®] 546 (sc-55518 AF546), Alexa Fluor[®] 594 (sc-55518 AF594) or Alexa Fluor[®] 647 (sc-55518 AF647), 200 μ g/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor[®] 680 (sc-55518 AF680) or Alexa Fluor[®] 790 (sc-55518 AF790), 200 μ g/ml, for Near-Infrared (NIR) WB, IF and FCM.

Blocking peptide available for competition studies, sc-55518 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

Alexa Fluor[®] is a trademark of Molecular Probes, Inc., Oregon, USA

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS


caveolin-3 (C-2) is recommended for detection of caveolin-3 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for caveolin-3 siRNA (h): sc-29943, caveolin-3 siRNA (m): sc-29944, caveolin-3 siRNA (r): sc-106997, caveolin-3 shRNA Plasmid (h): sc-29943-SH, caveolin-3 shRNA Plasmid (m): sc-29944-SH, caveolin-3 shRNA Plasmid (r): sc-106997-SH, caveolin-3 shRNA (h) Lentiviral Particles: sc-29943-V, caveolin-3 shRNA (m) Lentiviral Particles: sc-29944-V and caveolin-3 shRNA (r) Lentiviral Particles: sc-106997-V.


Molecular Weight of caveolin-3: 20-25 kDa.

Positive Controls: mouse heart extract: sc-2254, rat skeletal muscle extract: sc-364810 or rat heart extract: sc-2393.

DATA



caveolin-3 (C-2) HRP: sc-55518 HRP. Direct western blot analysis of caveolin-3 expression in mouse heart (A), rat skeletal muscle (B) and rat heart (C) tissue extracts.



caveolin-3 (C-2): sc-55518. Immunoperoxidase staining of formalin fixed, paraffin-embedded mouse heart muscle (A) and rat skeletal muscle (B) tissue showing membrane and cytoplasmic staining of myocytes.

SELECT PRODUCT CITATIONS

1. Ishii, M., et al. 2011. EphB signaling inhibits gap junctional intercellular communication and synchronized contraction in cultured cardiomyocytes. *Basic Res. Cardiol.* 106: 1057-1068.
2. Gómez-Díaz, B., et al. 2012. Immunodetection analysis of muscular dystrophies in Mexico. *Muscle Nerve* 45: 338-345.
3. Mitrofanova, L.B., et al. 2014. Evidence of specialized tissue in human interatrial septum: histological, immunohistochemical and ultrastructural findings. *PLoS ONE* 9: e113343.
4. Carmeille, R., et al. 2016. Membrane repair of human skeletal muscle cells requires Annexin-A5. *Biochim. Biophys. Acta* 1863: 2267-2279.
5. Hu, C.C., et al. 2022. Cardiac-targeted PIASy gene silencing mediates deSUMOylation of caveolin-3 and prevents ischemia/reperfusion-induced Na_v1.5 downregulation and ventricular arrhythmias. *Mil. Med. Res.* 9: 58.

RESEARCH USE

For research use only, not for use in diagnostic procedures.