

# Scrib (D-8): sc-55532

## BACKGROUND

*Drosophila melanogaster* genes, which are categorized based on the type of protein for which they encode, represent six major classifications, including intracellular signaling proteins, transmembrane proteins, RNA binding proteins, secreted factors, transcription regulators (basic helix-loop-helix, homeodomain containing, zinc finger containing and chromatin associated) and other functional proteins. Morphogenesis and cell differentiation in *Drosophila* requires accurate control of cell division. Discs large (Dlg), Scribble (Scrib) and Lethal giant larvae (Lgl) tumor suppressor proteins regulate multiple aspects of neuroblast asymmetric cell division. Dlg/Scrib/Lgl proteins show apical cortical enrichment at prophase/metaphase and have a uniform cortical distribution. Mutations in the genes encoding multi-PDZ (PSD-95, discs-large and ZO-1) and the leucine-rich-repeat protein Scrib cause aberrant cell shapes and the loss of monolayer organization of embryonic epithelia. The human homolog, hScrib, is intracellularly localized to the vertebrate tight junction, which functions to correctly place adherens junctions. The PDZ domains of Scrib are predicted to bind to the consensus S/TXV at the C-terminus of proteins. PDZ domain proteins have been implicated at several different sites of the protein trafficking pathway, suggesting that Scrib is required for the localization of several epithelial determinants.

## CHROMOSOMAL LOCATION

Genetic locus: SCRIB (human) mapping to 8q24.3.

## SOURCE

Scrib (D-8) is a mouse monoclonal antibody raised against amino acids 1331-1630 (deletion 1444-1504) mapping at the C-terminus of Scrib of human origin.

## PRODUCT

Each vial contains 200 µg IgM kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## APPLICATIONS

Scrib (D-8) is recommended for detection of Scrib of human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for Scrib siRNA (h): sc-36466, Scrib shRNA Plasmid (h): sc-36466-SH and Scrib shRNA (h) Lentiviral Particles: sc-36466-V.


Molecular Weight of Scrib: 210 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200, Hep G2 cell lysate: sc-2227 or Jurkat whole cell lysate: sc-2204.


## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein L-Agarose: sc-2336 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850. 4) Immunohistochemistry: use m-IgGκ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohisto-mount: sc-45086, or Organo/Limonene Mount: sc-45087.

## DATA



Scrib (D-8): sc-55532. Western blot analysis of Scrib expression in Hep G2 whole cell lysate.



Scrib (D-8): sc-55532. Immunoperoxidase staining of formalin fixed, paraffin-embedded human skin tissue showing cytoplasmic staining of epidermal cells.

## SELECT PRODUCT CITATIONS

- Vaira, V., et al. 2011. Aberrant overexpression of the cell polarity module scribble in human cancer. *Am. J. Pathol.* 178: 2478-2483.
- Kapil, S., et al. 2017. The cell polarity protein Scrib functions as a tumor suppressor in liver cancer. *Oncotarget* 8: 26515-26531.
- Abedrabbo, M. and Ravid, S. 2020. Scribble, Lgl1, and myosin II form a complex *in vivo* to promote directed cell migration. *Mol. Biol. Cell.* E-published.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.

## CONJUGATES

See **Scrib (C-6): sc-55543** for Scrib antibody conjugates, including AC, HRP, FITC, PE, and Alexa Fluor® 488, 546, 594, 647, 680 and 790.