

# IDH3B (M-17): sc-55674

## BACKGROUND

IDH3B (isocitrate dehydrogenase 3 NAD<sup>+</sup> β, NAD<sup>+</sup>-specific ICDH) is a 384 amino acid protein encoded by the human gene IDH3B. IDH3B belongs to the isocitrate and isopropylmalate dehydrogenases family and can bind one magnesium or manganese ion per subunit. It is usually found in the mitochondrion as a heterooligomer of subunits α, β, and γ in the apparent ratio of 2:1:1. Human NAD-dependent isocitrate dehydrogenase (IDH) is allosterically activated by ADP by lowering the K<sub>m</sub> for isocitrate. NAD-dependent isocitrate dehydrogenase is a tricarboxylic acid cycle enzyme that produces 2-oxoglutarate, an organic acid required by the glutamine synthetase/glutamate synthase cycle to assimilate ammonium.

## REFERENCES

- Kim, Y.O., et al. 1995. Characterization of a cDNA clone for human NAD<sup>+</sup>-specific isocitrate dehydrogenase α-subunit and structural comparison with its isoenzymes from different species. *Biochem. J.* 32: 295-296.
- Hong, G., et al. 1997. Molecular cloning of a highly conserved mouse and human integral membrane protein (Itn1) and genetic mapping to mouse chromosome 9. *Genomics* 31: 295-300.
- Huh, T.L., et al. 1997. Assignment of the human mitochondrial NAD<sup>+</sup>-specific isocitrate dehydrogenase α-subunit (IDH3A) gene to 15q25.1→25.2 by *in situ* hybridization. *Genomics* 32: 295-296.
- Dash, D.P., et al. 2006. Fine mapping of the keratoconus with cataract locus on chromosome 15q and candidate gene analysis. *Mol. Vis.* 12: 499-505.
- Soundar, S., et al. 2006. Identification of Mn<sup>2+</sup>-binding aspartates from α, β, and γ subunits of human NAD-dependent isocitrate dehydrogenase. *J. Biol. Chem.* 281: 21073-21081.
- Imabayashi, F., et al. 2006. Substrate-free structure of a monomeric NADP isocitrate dehydrogenase: an open conformation phylogenetic relationship of isocitrate dehydrogenase. *Proteins* 63: 100-112.
- Veena, C.K., et al. 2007. Mitochondrial dysfunction in an animal model of hyperoxaluria: A prophylactic approach with fucoidan. *Eur. J. Pharmacol.* 579: 330-336.

## CHROMOSOMAL LOCATION

Genetic locus: *Idh3b* (mouse) mapping to 2 F1.

## SOURCE

IDH3B (M-17) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the C-terminus of IDH3B of mouse origin.

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.

## PRODUCT

Each vial contains 200 μg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-55674 P, (100 μg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## APPLICATIONS

IDH3B (M-17) is recommended for detection of IDH3B of mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μg per 100-500 μg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

IDH3B (M-17) is also recommended for detection of IDH3B in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for IDH3B siRNA (m): sc-62492, IDH3B shRNA Plasmid (m): sc-62492-SH and IDH3B shRNA (m) Lentiviral Particles: sc-62492-V.


Molecular Weight of IDH3B: 42 kDa.

Positive Controls: c4 whole cell lysate: sc-364186 or mouse embryo extract: sc-364239.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## DATA



IDH3B (M-17): sc-55674. Western blot analysis of IDH3B expression in c4 whole cell lysate (A) and mouse embryo tissue extract (B).

## RESEARCH USE

For research use only, not for use in diagnostic procedures.