

IDH3G (C-19): sc-55677

BACKGROUND

IDH3G (isocitrate dehydrogenase [NAD] subunit γ (mitochondrial), NAD⁺-specific ICDH) is a 393 amino acid protein encoded by the human gene IDH3G. IDH3G belongs to the isocitrate and isopropylmalate dehydrogenases family and can bind one magnesium or manganese ion per subunit. It is usually found in the mitochondrion as a heterooligomer of subunits α , β , and γ in the apparent ratio of 2:1:1. Human NAD-dependent isocitrate dehydrogenase (IDH) is allosterically activated by ADP by lowering the K_m for isocitrate. NAD-dependent isocitrate dehydrogenase is a tricarboxylic acid cycle enzyme that produces 2-oxoglutarate, an organic acid required by the glutamine synthetase/glutamate synthase cycle to assimilate ammonium.

REFERENCES

- Kim, Y.O., et al. 1995. Characterization of a cDNA clone for human NAD⁺-specific isocitrate dehydrogenase α -subunit and structural comparison with its isoenzymes from different species. *Biochem. J.* 308: 63-68.
- Lee, P. and Colman, R.F. 2006. Thr 373, Asp 375, and Lys 260 are in the coenzyme site of porcine NADP-dependent isocitrate dehydrogenase. *Arch. Biochem. Biophys.* 450: 183-190.

CHROMOSOMAL LOCATION

Genetic locus: IDH3G (human) mapping to Xq28.

SOURCE

IDH3G (C-19) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the C-terminus of IDH3G of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-55677 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

IDH3G (C-19) is recommended for detection of IDH3G of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

IDH3G (C-19) is also recommended for detection of IDH3G in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for IDH3G siRNA (h): sc-62493, IDH3G shRNA Plasmid (h): sc-62493-SH and IDH3G shRNA (h) Lentiviral Particles: sc-62493-V.


Molecular Weight of IDH3G: 43 kDa.

Positive Controls: MCF7 whole cell lysate: sc-2206, SK-BR-3 cell lysate: sc-2218 or IDH3G (h): 293 Lysate: sc-110789.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



IDH3G (C-19): sc-55677. Western blot analysis of IDH3G expression in HT-1080 (A), RD (B), A-673 (C), MCF7 (D) and SK-BR-3 (E) whole cell lysates.



IDH3G (C-19): sc-55677. Western blot analysis of IDH3G expression in non-transfected 293: sc-110760 (A), human IDH3G transfected 293: sc-110789 (B), SK-BR-3 (C), HT-1080 (D), HeLa (E) and MCF7 (F) whole cell lysates.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

MONOS
Satisfaction
Guaranteed

Try **IDH3G (C-5): sc-365489** or **IDH3G (C-8): sc-398066**, our highly recommended monoclonal alternatives to IDH3G (C-19).