

GNL2 (C-16): sc-55687

BACKGROUND

GNL2 (autoantigen NGP-1, NOG2) is a nucleolar guanine-triphosphate binding protein that is ubiquitously expressed at low levels in almost all tissues. GNL2 is involved in the crucial process of trafficking proteins out of the nucleus. Specifically, it is a GTPase that interacts with the 60s preribosomal subunit in the nucleus and facilitates export of the subunit into the cytoplasm. GTPases are responsible for the hydrolysis of GTP by way of a protein region dubbed the G domain. GTPases are often involved in the translocating proteins through membranes gleaned energy for the activity by hydrolyzing GTP. GNL2 shares G domain homology and some functionality with nucleostemin (GNL3), another nuclear GTPase. Highest expression of GNL2 is found in testis.

CHROMOSOMAL LOCATION

Genetic locus: GNL2 (human) mapping to 1p34.3; Gnl2 (mouse) mapping to 4 D2.2.

SOURCE

GNL2 (C-16) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the C-terminus of GNL2 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-55687 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

GNL2 (C-16) is recommended for detection of GNL2 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

GNL2 (C-16) is also recommended for detection of GNL2 in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for GNL2 siRNA (h): sc-62685, GNL2 siRNA (m): sc-62686, GNL2 shRNA Plasmid (h): sc-62685-SH, GNL2 shRNA Plasmid (m): sc-62686-SH, GNL2 shRNA (h) Lentiviral Particles: sc-62685-V and GNL2 shRNA (m) Lentiviral Particles: sc-62686-V.


Molecular Weight of GNL2: 84 kDa.

Positive Controls: Jurkat whole cell lysate: sc-2204, Ramos cell lysate: sc-2216 or mouse testis extract: sc-2405.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



GNL2 (C-16): sc-55687. Western blot analysis of GNL2 expression in Jurkat whole cell lysate.



GNL2 (C-16): sc-55687. Immunofluorescence staining of methanol-fixed HeLa cells showing nuclear localization.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



Try **GNL2 (B-8): sc-514050** or **GNL2 (B-10): sc-376732**, our highly recommended monoclonal alternatives to GNL2 (C-16).