# EMP-1 (D-13): sc-55713



The Power to Question

#### **BACKGROUND**

Epithelial membrane protein-1 (EMP-1) is a four pass transmembrane protein consisting of 160 amino acids. It is a member of a small family of epithelial membrane proteins. EMP-1 is very similar in structure to its close relative, peripheral myelin protein 22 (PMP22). It is most predominantly expressed in tissues of the gastrointestinal tract but has also been found to be a junctional protein in the liver expressed along the intercellular border. EMP-1 directly interacts with the C-terminus of the P2X7 receptor and may be involved in membrane blebbing. EMP-1 may also be an important regulator in cell communication, signaling and adhesion. When EMP-1 is overexpressed, cell proliferation is inhibited, S phase is arrested and  $\rm G_1$  phase is prolonged in esophogeal cancer cells. EMP-1 may play a role in tumorigenesis and has been identified as a biomarker for gefitinib treatment resistance for patients with lung cancer.

#### **REFERENCES**

- Taylor, V., et al. 1995. Epithelial membrane protein-1, peripheral myelin protein 22 and lens membrane protein-20 define a novel gene family. J. Biol. Chem. 270: 28824-28833.
- 2. Chen, Y., et al. 1997. cDNA cloning, genomic structure, and chromosome mapping of the human epithelial membrane protein CL-20 gene (EMP1), a member of the PMP22 family. Genomics 41: 40-48.
- 3. Jetten, A.M., et al. 2000. The peripheral myelin protein 22 and epithelial membrane protein family. Prog. Nucleic Acid Res. Mol. Biol. 64: 97-129.
- 4. Wang, H.T., et al. 2002. Effect of EMP-1 gene on human esophageal cancer cell line. Ai Zheng 21: 229-232.
- Wilson, H.L., et al. 2002. Epithelial membrane proteins induce membrane blebbing and interact with the P2X7 receptor C-terminus. J. Biol. Chem. 277: 34017-34023.
- Liu, Y.H., et al. 2003. Differential expression of the epithelial membrane protein-1 of laryngeal carcinoma. Zhongguo Yi Xue Ke Xue Yuan Xue Bao 25: 47-51.
- Wang, H.T., et al. 2003. Analysis of gene expression profile induced by EMP-1 in esophageal cancer cells using cDNA microarray. World J. Gastroenterol. 9: 392-398.
- 8. Lee, H.S., et al. 2005. EMP-1 is a junctional protein in a liver stem cell line and in the liver. Biochem. Biophys. Res. Commun. 334: 996-1003.
- 9. Jain, A., et al. 2005. Epithelial membrane protein-1 is a biomarker of gefitinib resistance. Proc. Natl. Acad. Sci. USA 102:11858-11863.

## **CHROMOSOMAL LOCATION**

Genetic locus: EMP1 (human) mapping to 12p12.3; Emp1 (mouse) mapping to 6 G1.

## SOURCE

EMP-1 (D-13) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of EMP-1 of rat origin.

#### **PRODUCT**

Each vial contains 200  $\mu g$  IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-55713 P, (100  $\mu$ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

#### **APPLICATIONS**

EMP-1 (D-13) is recommended for detection of EMP-1 of rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

#### **RECOMMENDED SECONDARY REAGENTS**

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

#### **STORAGE**

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

### **RESEARCH USE**

For research use only, not for use in diagnostic procedures.

### **PROTOCOLS**

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

Santa Cruz Biotechnology, Inc. 1.800.457.3801 831.457.3801 fax 831.457.3801 Europe +00800 4573 8000 49 6221 4503 0 www.scbt.com