

PREP-2 (P-20): sc-55889

BACKGROUND

PREP-2 (Pbx-regulating protein-2), also known as PBX/knotted 1 homeobox 2 or PKNOX2, is a widely expressed protein belonging to the TALE (three amino acid loop extension)/MEIS family. PREP-2 is a DNA-binding protein that forms stable complexes with Pbx proteins. It is highly homologous to the related protein PREP-1, but displays a more restricted tissue distribution and a higher DNA-dissociation rate. Like PREP-1, PREP-2 forms a heterodimer with Pbx 1. The PREP-2-Pbx 1 dimer is relocated to the nucleus where it associates with HoxB1 to form a ternary complex. In contrast with PREP-1, which acts to increase transcriptional activation in this ternary complex, PREP-2 leads to a slight decrease in transcriptional activity of the ternary complex. Multiple isoforms exist for PREP-2, localizing to the nucleus or cytoplasm. Cytoplasmic isoforms are believed to colocalize with F-Actin, G-Actin and Tubulin/microtubules.

CHROMOSOMAL LOCATION

Genetic locus: PKNOX2 (human) mapping to 11q24.2; Pknx2 (mouse) mapping to 9 A4.

SOURCE

PREP-2 (P-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of PREP-2 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-55889 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

Available as TransCruz reagent for Gel Supershift and ChIP applications, sc-55889 X, 200 µg/0.1 ml.

APPLICATIONS

PREP-2 (P-20) is recommended for detection of PREP-2 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

PREP-2 (P-20) is also recommended for detection of PREP-2 in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for PREP-2 siRNA (h): sc-62856, PREP-2 siRNA (m): sc-62857, PREP-2 shRNA Plasmid (h): sc-62856-SH, PREP-2 shRNA Plasmid (m): sc-62857-SH, PREP-2 shRNA (h) Lentiviral Particles: sc-62856-V and PREP-2 shRNA (m) Lentiviral Particles: sc-62857-V.

PREP-2 (P-20) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.


Molecular Weight of PREP-2: 70 kDa.

Positive Controls: HeLa nuclear extract: sc-2120 or IMR-32 nuclear extract: sc-2148.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



PREP-2 (P-20): sc-55889. Western blot analysis of human recombinant PREP-2 fusion protein.

SELECT PRODUCT CITATIONS

- Coy, S., et al. 2011. A novel Gli3 enhancer controls the Gli3 spatiotemporal expression pattern through a TALE homeodomain protein binding site. *Mol. Cell. Biol.* 31: 1432-1443.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.


 MONOS
Satisfaction
Guaranteed

Try **PREP-2 (56.1): sc-101857**, our highly recommended monoclonal alternative to PREP-2 (P-20).