

GRASP (C-13): sc-55951

BACKGROUND

GRASP (GRP1-associated scaffold protein, tamalin) is a 395 amino acid protein encoded by the human gene GRASP. GRASP is a scaffold protein that comprises multiple protein-interacting domains, including a postsynaptic density protein (PSD-95)/discs-large/ZO-1 (PDZ) domain, a leucine-zipper region and a carboxyl-terminal PDZ-binding motif. GRASP is involved with intracellular trafficking and contributes to the macromolecular organization of group 1 metabotropic glutamate receptors (mGluRs) at synapses. GRASP forms a heteromer composed of GRASP, PSCD2 and at least one mGluR-1. It also interacts with PSCD3, mGluR-2, mGluR-3 and mGluR-5. GRASP is highly expressed in brain and has lower levels of expression in lung, heart, embryo, kidney and ovary.

REFERENCES

- Langille, S.E., Patki, V., Klarlund, J.K., Buxton, J.M., Holik, J.J., Chawla, A., Corvera, S. and Czech, M.P. 1999. ADP-ribosylation factor 6 as a target of guanine nucleotide exchange factor GRP1. *J. Biol. Chem.* 274: 27099-27104.
- Dominguez, O., Ashhab, Y., Sabater, L., Belloso, E., Caro, P. and Pujol-Borrell, R. 1999. Cloning of ARE-containing genes by AU-motif-directed display. *Genomics* 54: 278-286.
- Nevrivy, D.J., Peterson, V.J., Avram, D., Ishmael, J.E., Hansen, S.G., Dowell, P., Hruba, D.E., Dawson, M.I. and Leid, M. 2000. Interaction of GRASP, a protein encoded by a novel retinoic acid-induced gene, with members of the cytohesin family of guanine nucleotide exchange factors. *J. Biol. Chem.* 275: 16827-16836.
- Ye, D., Wei, M., McGuire, M., Huang, K., Kapadia, G., Herzberg, O., Martin, B.M. and Dunaway-Mariano, D. 2001. Investigation of the catalytic site within the ATP-grasp domain of *Clostridium symbiosum* pyruvate phosphate dikinase. *J. Biol. Chem.* 276: 37630-37639.
- Kitano, J., Kimura, K., Yamazaki, Y., Soda, T., Shigemoto, R., Nakajima, Y. and Nakanishi, S. 2002. Tamalin, a PDZ domain-containing protein, links a protein complex formation of group 1 metabotropic glutamate receptors and the guanine nucleotide exchange factor cytohesins. *J. Neurosci.* 22: 1280-1289.

CHROMOSOMAL LOCATION

Genetic locus: GRASP (human) mapping to 12q13.13; Grasp (mouse) mapping to 15 F2.

SOURCE

GRASP (C-13) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the C-terminus of GRASP of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-55951 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

GRASP (C-13) is recommended for detection of GRASP of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

GRASP (C-13) is also recommended for detection of GRASP in additional species, including porcine.

Suitable for use as control antibody for GRASP siRNA (h): sc-62423, GRASP siRNA (m): sc-62424, GRASP shRNA Plasmid (h): sc-62423-SH, GRASP shRNA Plasmid (m): sc-62424-SH, GRASP shRNA (h) Lentiviral Particles: sc-62423-V and GRASP shRNA (m) Lentiviral Particles: sc-62424-V.


Molecular Weight of GRASP: 43 kDa.

Positive Controls: mouse brain extract: sc-2253.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



GRASP (C-13): sc-55951. Western blot analysis of GRASP expression in mouse brain tissue extract.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.