

# Emt (SPM409): sc-56383

## BACKGROUND

The Tec family of non-receptor tyrosine kinases is composed of six proteins designated Tec, Emt (also known as Itk or Tsk), Btk (previously known as Atk, BPK or Emb), Bmx, Txk (also known as Rlk) and Dsrc28C. All members of the family contain SH3 and SH2 domains and, with the exception of Txk and Dsrc28C, also contain a pleckstrin homology (PH) and a Tec homology (TH) domain in their amino termini. Four alternatively spliced forms of Tec are found to be expressed broadly in cells Emt gene product associates with CD28 and becomes activated subsequent to CD28 ligation. Btk is necessary for proper B cell development, and mutations in the gene encoding Btk have been associated with families suffering from X-linked agammaglobulinemia, also referred to as Bruton's disease. The Bmx protein shares a high degree of homology with Btk and seems to be expressed at highest levels in the heart. Txk expression is T cell-specific, while expression of the *Drosophila* Tec homolog, Dsrc28C, is developmentally regulated.

## REFERENCES

1. Yamada, N., et al. 1993. Structure and expression of novel protein tyrosine kinases, Emb and Emt, in hematopoietic cells. *Biochem. Biophys. Res. Commun.* 192: 231-240.
2. Thomas, J.D., et al. 1993. Co-localization of X-linked agammaglobulinemia and X-linked immunodeficiency genes. *Science* 261: 355-358.
3. Tamagnone, L., et al. 1994. BMX, a novel nonreceptor tyrosine kinase gene of the BTK/ITK/TEC/TXK family located in chromosome Xp22.2. *Oncogene* 9: 3683-3688.
4. Haire, R.N., et al. 1994. Txk, a novel human tyrosine kinase expressed in T cells shares sequence identity with Tec family kinases and maps to 4p12. *Hum. Mol. Genet.* 3: 897-901.
5. August, A., et al. 1994. CD28 is associated with and induces the immediate tyrosine phosphorylation and activation of the Tec family kinase Itk/Emt in the human Jurkat leukemic T cell line. *Proc. Natl. Acad. Sci. USA* 91: 9347-9351.
6. Hu, Q., et al. 1995. Identification of Rlk, a novel protein tyrosine kinase with predominant expression in the T cell lineage. *J. Biol. Chem.* 270: 1928-1934.

## CHROMOSOMAL LOCATION

Genetic locus: ITK (human) mapping to 5q33.3; Itk (mouse) mapping to 11 B1.1.

## SOURCE

Emt (SPM409) is a mouse monoclonal antibody raised against recombinant N-terminal Emt of mouse origin.

## PRODUCT

Each vial contains 200 µg IgG<sub>1</sub> kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## APPLICATIONS

Emt (SPM409) is recommended for detection of Emt of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)] and kinase assay.

Suitable for use as control antibody for Emt siRNA (h): sc-35300, Emt siRNA (m): sc-35301, Emt shRNA Plasmid (h): sc-35300-SH, Emt shRNA Plasmid (m): sc-35301-SH, Emt shRNA (h) Lentiviral Particles: sc-35300-V and Emt shRNA (m) Lentiviral Particles: sc-35301-V.


Molecular Weight of Emt: 72 kDa.

Positive Controls: CCRF-CEM cell lysate: sc-2225, ALL-SIL whole cell lysate: sc-364356 or Jurkat whole cell lysate: sc-2204.


## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml).

## DATA



Emt (SPM409): sc-56383. Western blot analysis of Emt expression in CCRF-CEM (A) and ALL-SIL (B) whole cell lysates.



Emt (SPM409): sc-56383. Western blot analysis of Emt expression in Jurkat whole cell lysate.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.