

Gliadin (HYB 314-01): sc-57745

BACKGROUND

Gliadin is a glycoprotein that, along with glutenin, plays a role in the formation of gluten, an ergastic amorphous mixture of proteins found in wheat, rye and barley. Gluten is responsible for the elasticity of kneaded dough, which allows it to be leavened. Gliadin is approximately 60% soluble in ethanol and contains only intramolecule disulfide links. Gliadin is also found in a variety of foods as well as in beer, along with the glycoprotein hordein. Induction of zonulin release in intestinal epithelial cells is triggered by Gliadin. This causes an activation of the zonulin pathway by PKC-mediated cytoskeleton reorganization and tight junction opening leads to a rapid increase in intestinal permeability to macromolecules. Individuals with disorders such as celiac disease or Crohn's disease are sensitive to Gliadin since they lack the enzyme necessary for its digestion and can not tolerate it in their diet.

REFERENCES

1. Shearer, G., Patey, A.L. and MacWeeny, D.J. 1975. Wheat flour proteins: the selectivity of solvents and the stability of Gliadin and glutenin fractions of stored flours. *J. Sci. Food Agric.* 26: 337-344.
2. Matsushima, N., Creutz, C.E. and Kretsinger, R.H. 1990. Polyproline, β -turn helices. Novel secondary structures proposed for the tandem repeats within rhodopsin, synaptophysin, synexin, Gliadin, RNA polymerase II, hordein, and gluten. *Proteins* 7: 125-155.
3. Weber, P., Zimmer, K.P., Husemann, S. and Harms, E. 1998. False-positive Gliadin and endomysium antibodies and exocrine pancreatic insufficiency as pitfalls in the differential diagnosis of duodenal Crohn's versus celiac disease. *Am. J. Gastroenterol.* 93: 1177-1179.
4. Clemente, M.G., De Virgiliis, S., Kang, J.S., Macatagney, R., Musu, M.P., Di Pierro, M.R., Drago, S., Congia, M. and Fasano, A. 2003. Early effects of Gliadin on enterocyte intracellular signalling involved in intestinal barrier function. *Gut* 52: 218-23.
5. Monteiro, A.M., Arêas, E.P., Schröder, A. and Fa, N. 2004. Gliadin effect on fluctuation properties of phospholipid giant vesicles. *Colloids Surf. B, Biointerfaces* 34: 53-57.
6. Drago, S., El Asmar, R., Di Pierro, M., Grazia Clemente, M., Tripathi, A., Sapone, A., Thakar, M., Iacono, G., Carroccio, A., D'Agate, C., Not, T., Zampini, L., Catassi, C. and Fasano, A. 2006. Gliadin, zonulin and gut permeability: effects on celiac and non-celiac intestinal mucosa and intestinal cell lines. *Scand. J. Gastroenterol.* 41: 408-419.
7. Howdle, P.D. 2006. Gliadin, glutenin or both? The search for the Holy Grail in coeliac disease. *Eur. J. Gastroenterol. Hepatol.* 18: 703-706.
8. Orecchioni, A.M., Duclairoir, C., Renard, D. and Nakache, E. 2006. Gliadin characterization by SANS and Gliadin nanoparticle growth modelization. *J. Nanosci. Nanotechnol.* 6: 3171-3178.

SOURCE

Gliadin (HYB 314-01) is a mouse monoclonal antibody raised against a Gliadin-related peptide coupled to immunogenic carrier protein.

PRODUCT

Each vial contains 100 μ g IgG_{2a} in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

Gliadin (HYB 314-01) is recommended for detection of Gliadin of Common Wheat (*Triticum aestivum*) origin by solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

SELECT PRODUCT CITATIONS

1. Yuan, Y., Qiu, J., Lin, Z.T., Li, W., Haley, C., Mui, U.N., Ning, J., Tying, S.K. and Wu, T. 2019. Identification of novel autoantibodies associated with psoriatic arthritis. *Arthritis Rheumatol.* E-published.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.