Hep C NS3 (251): sc-57771



The Power to Question

BACKGROUND

The Hep C (Hepatitis C) is a small, enveloped, single-stranded, positive sense RNA virus belonging to the family *Flaviviridae*. Transmission of the virus occurs when blood from an infected individual enters the body of an uninfected individual. Hep C primarily replicates within hepatocytes in the liver, and circulating Hep C particles bind to receptors on the surface and enter these cells. Hep C replicates quickly, producing approximately one trillion particles each day in infected individuals. Hep C RNA polymerase has no proofreading function, so the virus has an exceptionally high mutation rate which may help it elude the host's immune system. Hep C infection results in chronic infections, liver cirrhosis and hepatocellular carcinoma in most people. Hep C NS3 (nonstructural protein 3) has both protease and helicase activity and is essential for Hep C replication and proliferation.

REFERENCES

- Watashi, K. and Shimotohno, K. 2003. The roles of Hepatitis C virus proteins in a novel action mechanism of the HCV core protein on gene regulation by nuclear hormone receptors. Cancer Sci. 94: 937-943.
- Acosta-Rivero, N., Rodriguez, A., Musacchio, A., Falcón, V., Suarez, V.M., Chavez, L., Morales-Grillo, J. and Duenas-Carrera, S. 2004. Nucleic acid binding properties and intermediates of HCV core protein multimerization in *Pichia pastoris*. Biochem. Biophys. Res. Commun. 323: 926-931.
- Sansonno, D., Lauletta, G. and Dammacco, F. 2004. Detection and quantitation of HCV core protein in single hepatocytes by means of laser capture microdissection and enzyme-linked immunosorbent assay. J. Viral Hepat. 11: 27-32.
- Umehara, T., Fukuda, K., Nishikawa, F., Sekiya, S., Kohara, M., Hasegawa, T. and Nishikawa, S. 2004. Designing and analysis of a potent bifunctional aptamers that inhibit protease and helicase activities of HCV NS3. Nucleic Acids Symp. Ser. 48: 195-196.
- 5. Alisi, A., Mele, R., Spaziani, A., Tavolaro, S., Palescandolo, E. and Balsano, C. 2005. Thr 446 phosphorylation of PKR by HCV core protein deregulates G_2/M phase HCC cells. J. Cell. Physiol. 205: 25-31.
- Carabaich, A., Ruvoletto, M., Bernardinello, E., Tono, N., Cavalletto, L., Chemello, L., Gatta, A. and Pontisso, P. 2005. Profiles of HCV core protein and viremia in chronic Hepatitis C: possible protective role of core antigen in liver damage. J. Med. Virol. 76: 55-60.
- 7. Gu, J., Wang, L., Che, Y., Liu, L., Jiang, L., Dong, S., Li, W. and Li, Q. 2005. Morphological alteration and biological properties of hepatocytes not related to tumorigenesis following transfection with HCV core protein. J. Viral Hepat. 12: 20-26.
- 8. Kimball, P., Verbeke, S. and Shiffman, M. 2005. HCV core protein augments cyclosporine immunosuppression. Transplant Proc. 37: 652-653.

SOURCE

Hep C NS3 (251) is a mouse monoclonal antibody raised against a chimeric polyprotein corresponding to 555 amino acids within the internal region of Hep $\rm C.$

PRODUCT

Each vial contains 100 $\mu g \ lgG_1$ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

Hep C NS3 (251) is recommended for detection of an epitope corresponding to amino acids 1252-1477 of the NS3 region of Hep C origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000); non cross-reactive with core, Envelope Protein M (EPM), NS3 or NS4 regions.

Molecular Weight of Hep C NS3: 70 kDa.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-lgG κ BP-HRP: sc-516102 or m-lgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker^{IM} Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

Santa Cruz Biotechnology, Inc. 1.800.457.3801 831.457.3801 Fax 831.457.3801 Europe +00800 4573 8000 49 6221 4503 0 www.scbt.com