

HHV-6 gp90 (300): sc-57806

BACKGROUND

The *Herpesviridae* family consists of DNA viruses that cause diseases in humans and other animals. This family is comprised of eight distinct viruses: HHV-1–HHV-8. Human herpes virus type 6 (HHV-6) and HHV-7 are associated with febrile illnesses and the childhood disease exanthem subitum, while HHV-8 resembles the Epstein-Barr virus in its possible transforming properties and may play a role in lymphomas and Kaposi's sarcoma. HHV-6, a newly described β -herpesvirus that shares homology with cytomegalovirus (CMV), consists of two closely related variants: HHV-6A and HHV-6B. HHV-6 infection is followed by persistence and latency in different tissues including monocytes/macrophages, salivary glands, brain and kidney. HHV-6 activation may play a role in the pathogenesis of certain demyelinating diseases such as progressive multifocal leukoencephalopathy (PML) and multiple sclerosis (MS). HHV-6 DNA is normally found as a marker of active viral infection in serum samples of MS patients.

REFERENCES

1. Ablashi, D.V., Zompetta, C., Lease, C., Josephs, S.F., Komaroff, A.L., Krueger, G.R., Henry, B., Balachandra, N., Lukau, J. and Salahuddin, S.Z. 1994. Human herpesvirus 6 (HHV-6) and chronic fatigue syndrome (CFS). *Can. Dis. Wkly. Rep.* 17: 33-40.
2. Jayavas, C., Balachandra, K., Wongchuee, S., Kositanont, U. and Warachit, P. 1997. The latency rate of human herpesvirus 6 (HHV-6) in positive and negative human immunodeficiency virus (HIV) infection of intravenous drug users (IVDU). *Asian Pac. J. Allergy Immunol.* 15: 29-33.
3. Levy, J.A. 1997. Three new human herpesviruses (HHV-6, -7 and -8). *Lancet* 349: 558-563.
4. Blumberg, B.M., Mock, D.J., Powers, J.M., Ito, M., Assouline, J.G., Baker, J.V., Chen, B. and Goodman, A.D. 2000. The HHV-6 paradox: ubiquitous commensal or insidious pathogen? A two-step *in situ* PCR approach. *J. Clin. Virol.* 16: 159-178.
5. Abdel-Haq, N.M. and Asmar, B.I. 2004. Human herpesvirus 6 (HHV-6) infection. *Indian J. Pediatr.* 71: 89-96.
6. Caserta, M.T., McDermott, M.P., Dewhurst, S., Schnabel, K., Carnahan, J.A., Gilbert, L., Lathan, G., Lofthus, G.K. and Hall, C.B. 2004. Human herpesvirus 6 (HHV-6) DNA persistence and reactivation in healthy children. *J. Pediatr.* 145: 478-484.
7. Hernández-Losa, J., Fedele, C.G., Tenorio, A., Fernández, V., Castellví, J., Pozo, F., Parada, C. and Ramón y Cajal, S. 2005. Lack of association of polyomavirus and herpesvirus types 6 and 7 in human lymphomas. *Cancer* 103: 293-298.
8. Merk, J., Schmid, F.X., Schwarz, S., Lehane, C., Boehm, S., Salzberger, B., Fleck, M. and Birnbaum, D.E. 2005. Fatal pulmonary failure attributable to viral pneumonia with human herpes virus 6 (HHV-6) in a young immunocompetent woman. *J. Intensive Care Med.* 20: 302-306.
9. Debarbieux, S., Deroo-Berger, M.C., Grande, S., Najioullah, F., Kanitakis, J., Faure, M. and Claudy, A. 2006. Drug hypersensitivity syndrome associated with a primary HHV-6 infection. *Ann. Dermatol. Venereol.* 133: 145-147.

SOURCE

HHV-6 gp90 (300) is a mouse monoclonal antibody raised against human herpesvirus type 6.

PRODUCT

Each vial contains 100 μ g IgG₂ in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

HHV-6 gp90 (300) is recommended for detection of a 90 kDa glycoprotein of HHV-6 origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1–2 μ g per 100–500 μ g of total protein (1 ml of cell lysate)], and immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500).

Molecular Weight of HHV-6 gp90: 90 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-mouse IgG-HRP: sc-2005 (dilution range: 1:2000-1:32,000) or Cruz Marker™ compatible goat anti-mouse IgG-HRP: sc-2031 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunohistochemistry: use ImmunoCruz™: sc-2050 or ABC: sc-2017 mouse IgG Staining Systems.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.