

HPV18 E6 (289-13965): sc-57835

BACKGROUND

Human papilloma viruses (HPVs) can be classified as either high risk or low risk according to their association with cancer. HPV16 and HPV18 are the most common of the high risk group, while HPV6 and HPV11 are among the low risk types. Approximately 90% of cervical cancers contain HPV DNA of the high risk types. Mutational analysis have shown that the E6 and E7 genes of the high risk HPVs are necessary and sufficient for HPV transforming function. The specific interactions of the E6 and E7 proteins with p53 and pRB, respectively, correlate with HPV high and low risk classifications. The high risk HPV E7 proteins bind to pRB with a higher affinity than do the low risk HPV proteins, and only the high risk HPV E6 proteins form detectable complexes with p53 *in vitro*.

REFERENCES

1. Reich, N.C., et al. 1983. Two distinct mechanisms regulate the levels of a cellular tumor antigen, p53. *Mol. Cell. Biol.* 3: 2143-2150.
2. zur Hausen, H. and Schneider, A. 1987. The role of papilloma viruses in human angongenital cancer. In Howley, P.M. and Salzman, N.P., eds., *The Papovaviridae, 2 Papillomaviruses*. New York: Plenum, 245-263.
3. Munger, K., et al. 1989. The E6 and E7 genes of the human papillomavirus type 16 together are necessary and sufficient for transformation of primary human keratinocytes. *J. Virol.* 63: 4417-4421.
4. Munger, K., et al. 1989. Complex formation of human papillomavirus E7 proteins with the retinoblastoma tumor suppressor gene product. *EMBO J.* 8: 4099-4105.
5. Hawley-Nelson, P., et al. 1989. HPV16 E6 and E7 proteins cooperate to immortalize human foreskin keratinocytes. *EMBO J.* 13: 3905-3910.
6. Riou, G., et al. 1990. Association between poor prognosis in early-stage invasive cervical carcinomas and non-detection of HPV DNA. *Lancet* 335: 1171-1174.
7. Werness, B.A., et al. 1990. Association of human papillomavirus types 16 and 18 E6 proteins with p53. *Science* 248: 76-79.
8. Huibregtse, J.M., et al. 1993. Cloning and expression of the cDNA for E6-AP, a protein that mediates the interaction of the human papillomavirus E6 oncoprotein with p53. *Mol. Cell. Biol.* 13: 775-784.

SOURCE

HPV18 E6 (289-13965) is a mouse monoclonal antibody raised against purified HPV18 E6- β -galactosidase fusion protein.

PRODUCT

Each vial contains 200 μ g IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

HPV18 E6 (289-13965) is recommended for detection of the early protein E6 of Human Papillomavirus (HPV) types 16 and 18 by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500).

Molecular Weight of HPV18 E6: 16 kDa.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850. 4) Immunohistochemistry: use m-IgG κ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

SELECT PRODUCT CITATIONS

1. Hsu, C.H., et al. 2012. The HPV E6 oncoprotein targets histone methyltransferases for modulating specific gene transcription. *Oncogene* 31: 2335-2349.
2. Zhao, L., et al. 2019. E6-induced selective translation of WNT4 and JIP2 promotes the progression of cervical cancer via a noncanonical WNT signaling pathway. *Signal Transduct. Target. Ther.* 4: 32.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.



See **HPV16 E6/18 E6 (C1P5): sc-460** for HPV16 E6/18 E6 antibody conjugates, including AC, HRP, FITC, PE, and Alexa Fluor[®] 488, 546, 594, 647, 680 and 790.