# SANTA CRUZ BIOTECHNOLOGY, INC.

# Phencyclidine (6E7): sc-57968



## BACKGROUND

Phencyclidine, a contraction of the chemical name phenylcyclohexylpiperidine, also abbreviated PCP, is a dissociative drug that was previously used as an anesthetic agent, exhibiting hallucinogenic and neurotoxic effects. It functions mainly as an agonist to the N-methyl-D-Aspartate (NMDA) receptor, causing prolonged depolarization of the neuron and effectively blocking the activity of the NMDA receptor. A schedule II drug, Phencyclidine may cause a certain type of brain damage, known as Olney's lesions. Phencyclidine has a molecular weight of 243.387 g/mol, and athough the main psychoactive effects of the drug only last hours, total elimination from the body is prolonged, usually extending over weeks.

## REFERENCES

- 1. Aguayo, L.G. and Albuquerque, E.X. 1988. Phencyclidine blocks two potassium currents in spinal neurons in cell culture. Brain Res. 436: 9-17.
- Anderson, A.J. and Harvey, A.L. 1988. Effects of the facilitatory compounds catechol, guanidine, noradrenaline and Phencyclidine on presynaptic currents of mouse motor nerve terminals. Naunyn Schmiedebergs Arch. Pharmacol. 338: 133-137.
- Ffrench-Mullen, J.M. and Rogawski, M.A. 1993. Phencyclidine block of calcium current in isolated guinea pig hippocampal neurones. J. Physiol. 456: 85-105.
- Corso, T.D., Sesma, M.A., Tenkova, T.I., Der, T.C., Wozniak, D.F., Farber, N.B. and Olney, J.W. 1997. Multifocal brain damage induced by Phencyclidine is augmented by pilocarpine. Brain Res. 752: 1-14.
- Sams-Dodd, F. 1999. Phencyclidine in the social interaction test: an animal model of schizophrenia with face and predictive validity. Rev. Neurosci. 10: 59-90.
- Wiley, J.L. and Compton, A.D. 2004. Progressive ratio performance following challenge with antipsychotics, amphetamine or NMDA antagonists in adult rats treated perinatally with Phencyclidine. Psychopharmacology 177: 170-177.
- Wang, C., Fridley, J. and Johnson, K.M. 2005. The role of NMDA receptor upregulation in Phencyclidine-induced cortical apoptosis in organotypic culture. Biochem. Pharmacol. 69: 1373-1383.
- Winter, J.C., Eckler, J.R., Rice, K.C. and Rabin, R.A. 2005. Serotonergic/ glutamatergic interactions: potentiation of Phencyclidine-induced stimulus control by citalopram. Pharmacol. Biochem. Behav. 81: 694-700.
- Enomoto, T., Noda, Y., Mouri, A., Shin, E.J., Wang, D., Murai, R., Hotta, K., Furukawa, H., Nitta, A., Kim, H.C. and Nabeshima, T. 2005. Long-lasting impairment of associative learning is correlated with a dysfunction of N-methyl-D-aspartate-extracellular signaling-regulated kinase signaling in mice after withdrawal from repeated administration of Phencyclidine. Mol. Pharmacol. 68: 1765-1774.

#### SOURCE

Phencyclidine (6E7) is a mouse monoclonal antibody raised against Phencyclidine conjugated to BSA.

## PRODUCT

Each vial contains 100  $\mu g$  lgG\_1 kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

#### **APPLICATIONS**

Phencyclidine (6E7) is recommended for detection of Phencyclidine by solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

#### **RESEARCH USE**

For research use only, not for use in diagnostic procedures.

#### PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.