



## Salmonella (6321): sc-58018

### BACKGROUND

*Salmonella* bacteria are the most frequently reported cause of foodborne illness. *Salmonella* is a genus composed of rod-shaped, highly mobile Gram-negative bacterium. This non spore-forming genus includes over 2,000 serotypes of *Salmonella* bacteria, organized into five different serogroups: *Salmonella* A, B, C, D and E. A widespread occurrence of *Salmonella* in animals, chiefly in swine and poultry, and also environmental sources, including water, soil, insects, kitchen surfaces, factory surfaces, raw meats and animal feces, causes Salmonellosis. Several species are mildly pathogenic, producing slight gastroenteritis, while others generate a case of serious and often fatal food poisoning. Symptoms are reminiscent of the flu and may include myalgia, fever, lethargy, cough, gastrointestinal discomfort and sore throat. There is no difference in taste, smell or appearance in food infected with the bacteria. The bacteria persist in the intestinal tracts of infected animals and humans.

### REFERENCES

- Goh, Y.L., Yasin, R., Puthuchery, S.D., Koh, Y.T., Lim, V.K., Taib, Z. and Thong, K.L. 2003. DNA fingerprinting of human isolates of *Salmonella enterica* serotype Par B in Malaysia. *J. Appl. Microbiol.* 95: 1134-1142.
- Kudalkar, D., Thermidor, M. and Cunha, B.A. 2004. *Salmonella paratyphi* A enteric fever mimicking viral meningitis. *Heart Lung* 33: 414-416.
- Van Immerseel, F., Meulemans, L., De Buck, J., Pasmans, F., Velge, P., Bottreau, E., Haesebrouck, F. and Ducatelle, R. 2004. Bacteria-host interactions of *Salmonella Paratyphi* B dT+ in poultry. *Epidemiol. Infect.* 132: 239-243.
- Adachi, T., Sagara, H., Hirose, K. and Watanabe, H. 2005. Fluoroquinolone-resistant *Salmonella Paratyphi* A. *Emerg. Infect. Dis.* 11: 172-174.
- Erdem, B., Ercis, S., Hascelik, G., Gur, D., Gedikoglu, S., Aysev, A.D., Sumerkan, B., Tatman-Otkun, M. and Tuncer, I. 2005. Antimicrobial resistance patterns and serotype distribution among *Salmonella enterica* strains in Turkey, 2000-2002. *Eur. J. Clin. Microbiol. Infect. Dis.* 24: 220-225.
- Itah, A.Y. and Uweh, E.E. 2005. Bacteria isolated from blood, stool and urine of typhoid patients in a developing country. *Southeast Asian J. Trop. Med. Public Health* 36: 673-677.
- Khan, F.Y., Kamha, A.A. and Alomary, I.Y. 2006. Fulminant hepatic failure caused by *Salmonella paratyphi* A infection. *World J. Gastroenterol.* 12: 5253-5255.
- Mao, Y.F., Lin, X.J., Li, J., Ruan, P., Zhou, X.H. and Yan, J. 2006. Construction of prokaryotic expression system of *Salmonella paratyphi* A *spaO* gene and immunogenicity and immunoprotection of the expressed product. *Zhonghua Liu Xing Bing Xue Za Zhi* 27: 347-350.
- Morgan, J.S. and Laber, D.A. 2006. Ceftriaxone-resistant *Salmonella septicemia* and osteomyelitis in sickle cell disease adults. *South. Med. J.* 99: 625-627.

### SOURCE

Salmonella (6321) is a mouse monoclonal antibody raised against a *Salmonella* cell preparation.

### PRODUCT

Each vial contains 100 µg IgG<sub>2a</sub> in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

### APPLICATIONS

Salmonella (6321) is recommended for detection of flagellum protein present in most *Salmonella* species by solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

### STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

### RESEARCH USE

For research use only, not for use in diagnostic procedures.

### PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.