



# Yersinia pestis (6031): sc-58084

## BACKGROUND

*Yersinia pestis* is a Gram-negative coccobacillus belonging to the family *Enterobacteriaceae*. *Y. pestis* is primarily a rodent pathogen, with humans being an accidental host when bitten by an infected rat flea. It has a number of virulence factors that enable it to survive in humans by facilitating use of host nutrients, causing damage to host cells and subverting phagocytosis and other host defense mechanisms. The plasmid-encoded protein, virulence antigen (v), is a major protective immunogen that is involved in the translocation of the collection of toxins called *Yersinia* outer proteins (YOPs). The transcriptional activator PhoP is essential for survival of *Yersinia pestis* in macrophage phagosomes. However, the phagosomes occupied by *Y. pestis* have not been well characterized, and the mechanism by which PhoP promotes bacterial survival in these vacuoles is not fully understood.

## REFERENCES

1. Ferreras, J.A., Ryu, J.S., Di Lello, F., Tan, D.S. and Quadri, L.E. 2005. Small-molecule inhibition of siderophore biosynthesis in *Mycobacterium tuberculosis* and *Yersinia pestis*. *Nat. Chem. Biol.* 1: 29-32.
2. Honko, A.N., Sriranganathan, N., Lees, C.J. and Mizel, S.B. 2006. Flagellin is an effective adjuvant for immunization against lethal respiratory challenge with *Yersinia pestis*. *Infect. Immun.* 74: 1113-1120.
3. Parent, M.A., Wilhelm, L.B., Kummer, L.W., Szaba, F.M., Mullarky, I.K. and Smiley, S.T. 2006.  $\gamma$  interferon, tumor necrosis factor  $\alpha$ , and nitric oxide synthase 2, key elements of cellular immunity, perform critical protective functions during humoral defense against lethal pulmonary *Yersinia pestis* infection. *Infect. Immun.* 74: 3381-3386.
4. Grabenstein, J.P., Fukuto, H.S., Palmer, L.E. and Bliska, J.B. 2006. Characterization of phagosome trafficking and identification of PhoP-regulated genes important for survival of *Yersinia pestis* in macrophages. *Infect. Immun.* 74: 3727-3741.
5. Khushiramani, R., Shukla, J., Tuteja, U. and Batra, H.V. 2006. *Yersinia* outer-membrane protein B (YopB): a tool for identification of *Yersinia pestis* isolates. *J. Med. Microbiol.* 55: 467-469.
6. Khushiramani, R., Tuteja, U., Shukla, J., Panikkar, A. and Batra, H.V. 2006. Virulence markers of LCR plasmid in Indian isolates of *Yersinia pestis*. *APMIS* 114: 15-22.
7. Tan, L. and Darby, C. 2006. *Yersinia pestis* YrbH is a multifunctional protein required for both 3-deoxy-D-manno-oct-2-ulosonic acid biosynthesis and biofilm formation. *Mol. Microbiol.* 61: 861-870.
8. Torrea, G., Chenal-Francisque, V., Leclercq, A. and Carniel, E. 2006. Efficient tracing of global isolates of *Yersinia pestis* by restriction fragment length polymorphism analysis using three insertion sequences as probes. *J. Clin. Microbiol.* 44: 2084-2092.

## SOURCE

*Yersinia pestis* (6031) is a mouse monoclonal antibody raised against a *Yersinia pestis* preparation.

## PRODUCT

Each vial contains 100  $\mu$ g IgG<sub>1</sub> in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

## APPLICATIONS

*Yersinia pestis* (6031) is recommended for detection of *Yersinia pestis* by immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

Molecular Weight of *Yersinia pestis*: 21 kDa.

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.