

Diphtheria Toxin (7801): sc-58119

BACKGROUND

Corynebacterium diphtheriae is a Gram-positive, nonmotile aerobic bacteria found in soil and animal feces. *C. diphtheriae* bacteria infect the epithelial cells of the upper respiratory tract and, from there, produce and secrete a potent toxin which is absorbed and disseminated through lymph channels and blood to the susceptible tissues of the body. Diphtheria Toxin catalyzes the ADP-ribosylation and inactivation of eEF-2. The structure of Diphtheria Toxin reveals a Y-shaped molecule of three domains: a catalytic domain (fragment A), whose fold is of the $\alpha + \beta$ type; a transmembrane (TM) domain consisting of nine α -helices, two pairs of which may participate in pH-triggered membrane insertion and translocation; and a receptor-binding domain, which forms a flattened β -barrel with a jelly-roll-like topology. Together the TM- and receptor binding-domains constitute fragment B.

REFERENCES

1. Davis, L., et al. 1919. Studies on diphtheria toxin: II. The role of the amino acids in the metabolism of *Bacterium diphtheriae*. *J. Bacteriol.* 4: 217-241.
2. Hazen, E.L., et al. 1932. Further studies upon the effect of various carbohydrates on production of diphtheria toxin with special reference to its flocculating titer and final pH. *J. Bacteriol.* 23: 195-209.
3. Moloney, P.J., et al. 1932. The effect on solutions of diphtheria toxin and diphtheria antitoxin of contact with certain surfaces. *Biochem. J.* 26: 1754-1761.
4. Eaton, M.D. 1936. The purification and concentration of diphtheria toxin: II. Observations on the nature of the toxin. *J. Bacteriol.* 31: 367-383.
5. Eaton, M.D. 1937. The purification and concentration of diphtheria toxin: III. Separation of toxin from bacterial protein. *J. Bacteriol.* 34: 139-151.
6. Pappenheimer, A.M. 1942. Studies on diphtheria toxin and its reaction with antitoxin. *J. Bacteriol.* 43: 273-289.
7. Collier, R.J. 1975. Diphtheria toxin: mode of action and structure. *Bacteriol. Rev.* 39: 54-85.
8. Pappenheimer, A.M. 1977. Diphtheria toxin. *Annu. Rev. Biochem.* 46: 69-94.
9. Donovan, J.J., et al. 1981. Diphtheria toxin forms transmembrane channels in planar lipid bilayers. *Proc. Natl. Acad. Sci. USA* 78: 172-176.

SOURCE

Diphtheria Toxin (7801) is a mouse monoclonal antibody raised against Diphtheria Toxin.

PRODUCT

Each vial contains 100 μ g IgG₁ in 1.0 ml PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

Diphtheria Toxin (7801) is recommended for detection of different determinants of Diphtheria Toxin and anatoxin of *Corynebacterium diphtheriae* origin by solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.