



Toxoplasma gondii (6201): sc-58202

BACKGROUND

Toxoplasma gondii, a single-celled parasite, causes the disease toxoplasmosis. The life cycle of *Toxoplasma gondii* includes two phases, called the intestinal (or enteroepithelial) and extraintestinal phases. The intestinal phase produces oocysts and occurs only in wild and domesticated cats. The extraintestinal phase occurs in all infected animals and produces tachyzoites (actively proliferating trophozoites) and, eventually, bradyzoites (slowly growing trophozoites) or zoitocysts. *Toxoplasma gondii* infects tissue of the gastrointestinal tract; an active infection is accompanied by fever and enlargement of the spleen. Transmission of *Toxoplasma gondii* from pregnant women to their fetus can result in severe consequences.

REFERENCES

1. Dubey, J.P. 1998. Advances in the life cycle of *Toxoplasma gondii*. Int. J. Parasitol. 28: 1019-1024.
2. Dubey, J.P., Lindsay, D.S. and Speer, C.A. 1998. Structures of *Toxoplasma gondii* tachyzoites, bradyzoites and sporozoites and biology and development of tissue cysts. Clin. Microbiol. Rev. 11: 267-299.
3. Bonfioli, A.A. and Orefice, F. 2005. Toxoplasmosis. Semin. Ophthalmol. 20: 129-141.
4. Kravetz, J.D. and Federman, D.G. 2005. Toxoplasmosis in pregnancy. Am. J. Med. 118: 212-216.
5. Lüder, C.G. and Gross, U. 2005. Apoptosis and its modulation during infection mechanisms and role in pathogenesis. Curr. Top. Microbiol. Immunol. 289: 219-237.
6. Simpson, K.E., Devine, B.C. and Gunn-Moore, D. 2005. Suspected *Toxoplasma*-associated myocarditis in a cat. J. Feline Med. Surg. 7: 203-208.
7. Kim, J.Y., Ahn, M.H., Song, H.O., Choi, J.H., Ryu, J.S., Min, D.Y. and Cho, M.H. 2006. Involvement of MAPK activation in chemokine or COX-2 productions by *Toxoplasma gondii*. Korean J. Parasitol. 44: 197-207.
8. Mazumdar, J., H Wilson, E., Masek, K., A Hunter, C. and Striepen, B. 2006. Apicoplast fatty acid synthesis is essential for organelle biogenesis and parasite survival in *Toxoplasma gondii*. Proc. Natl. Acad. Sci. USA 103: 13192-13197.
9. Rorman, E., Zamir, C.S., Rilakis, I. and Ben-David, H. 2006. Congenital toxoplasmosis—prenatal aspects of *Toxoplasma gondii* infection. Reprod. Toxicol. 21: 458-472.

SOURCE

Toxoplasma gondii (6201) is a mouse monoclonal antibody raised against *Toxoplasma gondii*.

PRODUCT

Each vial contains 100 µg IgG_{2a} in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

Toxoplasma gondii (6201) is recommended for detection of *Toxoplasma gondii* origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) and immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

Molecular Weight of Toxoplasma gondii: 36 kDa.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.