

cathepsin W (BV39-2B): sc-58337

BACKGROUND

Cathepsin W (lymphopain) and cathepsin F comprise a novel subgroup of cathepsin proteases, and are phylogenetically distinct from other human cathepsins. The cathepsin W gene maps to chromosome 11q13.1 and contains ten exons with introns ranging from 81-119 bp. Cathepsin W protein is expressed specifically in CD8⁺ T lymphocytes. The expression of cathepsin W first occurs during the differentiation of thymocytes to CD8⁺ T lymphocytes, just as the thymocytes cease expression of CD4⁺ receptors. In transfected Cos-7 and HeLa cells, cathepsin W localizes within the rough endoplasmic reticulum. Cathepsin W contains a unique 21 amino acid peptide insertion between the active site histidine and asparagine residues, in addition to a distinctive 8-amino acid carboxy-terminal extension. An extended loop structure in the second or beta-sheet domain and an additional disulfide bond are two of several signature features of cathepsin W. Other features of cathepsin W include an additional cysteine, an S2 pocket and an additional residue. Cathepsin W may exist as a dimer with each monomer forming a disulfide bond.

REFERENCES

- Linnevers, C., et al. 1997. Human cathepsin W, a putative cysteine protease predominantly expressed in CD8⁺ T lymphocytes. *FEBS Lett.* 405: 253-259.
- Wex, T., et al. 1998. Genomic structure, chromosomal localization and expression of human cathepsin W. *Biochem. Biophys. Res. Commun.* 248: 255-261.
- Wex, T., et al. 1999. Human cathepsins F and W: a new subgroup of cathepsins. *Biochem. Biophys. Res. Commun.* 259: 401-407.
- Bhandoola, A., et al. 2000. Programming for cytotoxic effector function occurs concomitantly with CD4 extinction during CD8⁺ T cell differentiation in the thymus. *Int. Immunol.* 12: 1035-1040.
- Brinkworth, R.I., et al. 2000. Phylogenetic relationships and theoretical model of human cathepsin W (lymphopain), a cysteine proteinase from cytotoxic T lymphocytes. *Int. J. Biochem. Cell Biol.* 32: 373-384.

CHROMOSOMAL LOCATION

Genetic locus: CTSW (human) mapping to 11q13.1.

SOURCE

cathepsin W (BV39-2B) is a mouse monoclonal antibody raised against recombinant procathepsin W of human origin.

PRODUCT

Each vial contains 100 µg IgG_{2b} in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS


cathepsin W (BV39-2B) is recommended for detection of cathepsin W and procathepsin W of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)] and immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

Suitable for use as control antibody for cathepsin W siRNA (h): sc-72807, cathepsin W shRNA Plasmid (h): sc-72807-SH and cathepsin W shRNA (h) Lentiviral Particles: sc-72807-V.

Molecular Weight of cathepsin W: 47 kDa.

Positive Controls: Jurkat whole cell lysate: sc-2204 or cathepsin W (h): 293T Lysate: sc-116060.

DATA



cathepsin W (BV39-2B): sc-58337. Western blot analysis of cathepsin W expression in non-transfected: sc-117752 (A) and human cathepsin W transfected: sc-116060 (B) 293T whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.