

Rad54B (N-16): sc-5852

BACKGROUND

Rad52 family members (Rad50, Rad51B/C/D, Rad52, Rad54, MRE11) mediate DNA double-strand break repair (DSBR) for DNA damage that otherwise could cause cell death, mutation or neoplastic transformation. Rad51 (RECA, BRCC5) interacts with BRCA1 and BRCA2 to influence subcellular localization and cellular response to DNA damage. BRCA2 inactivation may be a key event leading to genomic instability and tumorigenesis from deregulation of Rad51. Rad52 forms a heptameric ring that binds single-stranded DNA ends and catalyzes DNA-DNA interaction necessary for the annealing of complementary strands. Rad52 can interact with Rad51. Rad54A of the DEAD-like helicase superfamily binds to double-strand DNA and induces a DNA topological change, which is thought to facilitate homologous DNA pairing and stimulate DNA recombination. Rad54B of the DEAD-like helicase superfamily binds to double-stranded DNA and displays ATPase activity in the presence of DNA. RAD54B is abundant in testis and spleen, and mutations of this gene occur in primary lymphoma and colon cancer. MRE11 (meiotic recombination 11, ATLD, HNGS1) is a nuclear 3'-5' exonuclease/endonuclease that associates with RAD50 and influences homologous recombination, telomere length maintenance, and DNA double-strand break repair. MRE11 is most abundant in proliferating tissues.

CHROMOSOMAL LOCATION

Genetic locus: RAD54B (human) mapping to 8q22.1; E130016E03Rik (mouse) mapping to 4 A1.

SOURCE

Rad54B (N-16) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of Rad54B of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.
 Blocking peptide available for competition studies, sc-5852 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

Rad54B (N-16) is recommended for detection of Rad54B of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Rad54B (N-16) is also recommended for detection of Rad54B in additional species, including equine.


Suitable for use as control antibody for Rad54B siRNA (h): sc-37401, Rad54B siRNA (m): sc-37402, Rad54B shRNA Plasmid (h): sc-37401-SH, Rad54B shRNA Plasmid (m): sc-37402-SH, Rad54B shRNA (h) Lentiviral Particles: sc-37401-V and Rad54B shRNA (m) Lentiviral Particles: sc-37402-V.

Molecular Weight of soluble Rad54B: 100 kDa.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



Rad54B (N-16): sc-5852. Western blot analysis of Rad54B expression in non-transfected 293: sc-110760 (A), human Rad54B transfected 293: sc-110613 (B), CCRF-CEM (C) and DU 145 (D) whole cell lysates and HeLa nuclear extract (E).



Rad54B (N-16): sc-5852. Western blot analysis of Rad54B expression in UV-irradiated HeLa cells.

STORAGE

Store at 4° C, **DO NOT FREEZE** Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



Try **Rad54B (19-K2): sc-101234**, our highly recommended monoclonal alternative to Rad54B (N-16).