SANTA CRUZ BIOTECHNOLOGY, INC.

UCH-L1 (31A3): sc-58595



BACKGROUND

UCH-L1 (ubiquitin C-terminal hydrolase) is a member of a gene family whose products hydrolyze small C-terminal adducts of ubiquitin to generate the ubiquitin monomer. Expression of UCH-L1 is highly specific to neurons and to cells of the diffuse neuroendocrine system and their tumors. UCH-L1 is expressed in brain neurons. Examination of specific brain regions reveals expression in all areas tested, particularly in the substantia nigra. UCH-L1 represents 1 to 2% of total soluble brain protein. Its occurrence in Lewy bodies and its function in the proteasome pathway make it a compelling candidate gene in Parkinson disease. The gene which encodes UCH-L1 maps to human chromosome 4p13. The 230 amino acid human UCH-L3 protein is 54% identical to that of UCH-L1. UCH-L3 is the predominant thiol protease and has high-affinity binding sites for ubiquitin.

CHROMOSOMAL LOCATION

Genetic locus: UCHL1 (human) mapping to 4p13; Uchl1 (mouse) mapping to 5 C3.1.

SOURCE

UCH-L1 (31A3) is a mouse monoclonal antibody raised against native UCH-L1 of human origin.

PRODUCT

Each vial contains 200 μ g lgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

UCH-L1 (31A3) is recommended for detection of UCH-L1 on neuronal cell bodies and axons in the CNS and periphery, small nerve fibers in peripheral tissues, neuroendocrine cells in the pituitary, thyroid, pancreas and tumors of the DNES of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500).

UCH-L1 (31A3) is also recommended for detection of UCH-L1 on neuronal cell bodies and axons in the CNS and periphery, small nerve fibers in peripheral tissues, neuroendocrine cells in the pituitary, thyroid, pancreas and tumors of the DNES in additional species, including bovine and porcine.

Suitable for use as control antibody for UCH-L1 siRNA (h): sc-42304, UCH-L1 siRNA (m): sc-42305, UCH-L1 shRNA Plasmid (h): sc-42304-SH, UCH-L1 shRNA Plasmid (m): sc-42305-SH, UCH-L1 shRNA (h) Lentiviral Particles: sc-42304-V and UCH-L1 shRNA (m) Lentiviral Particles: sc-42305-V.

Molecular Weight of UCH-L1: 25 kDa.

Positive Controls: SK-N-SH cell lysate: sc-2410, rat brain extract: sc-2392 or SH-SY5Y cell lysate: sc-3812.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850. 4) Immunohistochemistry: use m-IgGκ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

DATA





UCH-L1 (31A3): sc-58595. Western blot analysis of UCH-L1 expression in SH-SYSY (A), SK-N-SH (B), NCI-H292 (C), RIN-m5F (D), H19-7/IGF-IR (E) and PC-12 (F) whole cell lysates.

UCH-L1 expression in rat brain tissue extract.

SELECT PRODUCT CITATIONS

- Yoshida, A., et al. 2012. *In vitro* tissue engineering of smooth muscle sheets with peristalsis using a murine induced pluripotent stem cell line. J. Pediatr. Surg. 47: 329-335.
- 2. Lee, J.H., et al. 2023. Stem cell properties of gastric cancer stem-like cells under stress conditions are regulated via the c-Fos/UCH-L3/ β -catenin axis. Mol. Cells 46: 476-485.

STORAGE

Store at 4° C, **D0 NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.