


CD163 (ED2): sc-58965



The Power to Question

BACKGROUND

CD163, also designated M130, is a macrophage-associated antigen that is a member of the scavenger receptor cysteine-rich (SRCR) superfamily. It is highly expressed on macrophages and to a lesser extent on monocytes. The acute phase-regulated and signal-inducing macrophage protein, CD163, is a receptor that scavenges hemoglobin by mediating endocytosis of haptoglobin-hemoglobin complexes. CD163 binds only haptoglobin and hemoglobin in complex, which indicates the exposure of a receptor-binding neopeptide. The receptor-ligand interaction is calcium-dependent and of high affinity. The existence of several CD163 isoforms, which differ in the structure of their cytoplasmic domains and putative phosphorylation sites, suggests that these isoforms also differ in their signaling mechanism. The gene which encodes CD163 maps to human chromosome 12p13.31.

SOURCE

CD163 (ED2) is a mouse monoclonal antibody raised against a spleen cell homogenate of rat origin.

PRODUCT

Each vial contains 100 µg IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

CD163 (ED2) is recommended for detection of the CD163 cell surface glycoprotein of mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and flow cytometry (1 µg per 1 x 10⁶ cells).

Molecular Weight of CD163: 130 kDa.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850. 4) Immunohistochemistry: use m-IgGκ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

SELECT PRODUCT CITATIONS

- Omar, O., et al. 2010. *In vivo* gene expression in response to anodically oxidized versus machined titanium implants. *J. Biomed. Mater. Res. A* 92: 1552-1566.
- Omar, O., et al. 2010. Integrin and chemokine receptor gene expression in implant-adherent cells during early osseointegration. *J. Mater. Sci. Mater. Med.* 21: 969-980.
- Vaughn, C.N., et al. 2013. Cellular neuroinflammation in a lateral forceps compression model of spinal cord injury. *Anat. Rec.* 296: 1229-1246.
- Yuan, S., et al. 2014. Effect of bone marrow mesenchymal stem cell transplantation on acute hepatic failure in rats. *Exp. Ther. Med.* 8: 1150-1158.
- Zhang, X.L., et al. 2014. Vitamin D prevents podocyte injury via regulation of macrophage M₁/M₂ phenotype in diabetic nephropathy rats. *Endocrinology* 155: 4939-4950.
- Sawano, T., et al. 2015. Fatty acid synthase-positive hepatocytes and subsequent steatosis in rat livers by irinotecan. *Oncol. Rep.* 33: 2151-2160.
- Alan, E., et al. 2015. The profile of the epidermal growth factor system in rat endometrium during postpartum involution period. *Vet. Res. Commun.* 39: 115-135.
- Suciu, M., et al. 2015. Acetaminophen-induced liver injury: Implications for temporal homeostasis of lipid metabolism and eicosanoid signaling pathway. *Chem. Biol. Interact.* 242: 335-344.
- Elgali, I., et al. 2016. Guided bone regeneration using resorbable membrane and different bone substitutes: Early histological and molecular events. *Acta Biomater.* 29: 409-423.
- Lu, W., et al. 2016. The new role of CD163 in the differentiation of bone marrow stromal cells into vascular endothelial-like cells. *Stem Cells Int.* 2016: 2539781.
- Li, Y., et al. 2016. Gelatin microgel incorporated poly(ethylene glycol)-based bioadhesive with enhanced adhesive property and bioactivity. *ACS Appl. Mater. Interfaces* 8: 11980-11989.
- D'Amore, A., et al. 2016. Bi-layered polyurethane-extracellular matrix cardiac patch improves ischemic ventricular wall remodeling in a rat model. *Biomaterials* 107: 1-14.
- Lee, R.S., et al. 2017. The influence of titanium surface characteristics on macrophage phenotype polarization during osseous healing in type I diabetic rats: a pilot study. *Clin. Oral Implants Res.* 28: e159-e168.
- Meng, H., et al. 2017. Model polymer system for investigating the generation of hydrogen peroxide and its biological responses during the crosslinking of mussel adhesive moiety. *Acta Biomater.* 48: 144-156.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.