

SSRP1 (D-15): sc-5909

BACKGROUND

Expression of protein-coding genes requires the association of specific transcription factors, RNA polymerase and various accessory factors. These accessory factors are distinguished as either histone acetyltransferases or ATP-dependent chromatin-remodeling enzymes, which include FACT (for facilitates chromatin transcription), and they facilitate transcription initiation on DNA packaged into chromatin. FACT is a chromatin-specific elongation factor required for transcription of chromatin templates, and it specifically interacts with nucleosomes and histone H2A/H2B dimers, to promote nucleosome disassembly upon transcription. FACT represents a complex between SPT16, a homologue of the *Saccharomyces cerevisiae* Spt16/Cdc68 protein, and the high-mobility group (HMG)-1-like protein structure-specific recognition protein-1 (SSRP-1). Similar to other (HMG) domain containing proteins, which are characterized by their ability to bend target DNAs, SSRP1 and the murine ortholog T160 physically interact with serum response factors (SRF) and function as positive co-regulatory proteins involved in modulating SRF-dependent gene expression.

CHROMOSOMAL LOCATION

Genetic locus: SSRP1 (human) mapping to 11q12; Ssrp1 (mouse) mapping to 2 D.

SOURCE

SSRP1 (D-15) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of SSRP1 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-5909 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

SSRP1 (D-15) is recommended for detection of SSRP1 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

SSRP1 (D-15) is also recommended for detection of SSRP1 in additional species, including canine, bovine, porcine and avian.

Suitable for use as control antibody for SSRP1 siRNA (h): sc-37877, SSRP1 siRNA (m): sc-37878, SSRP1 shRNA Plasmid (h): sc-37877-SH, SSRP1 shRNA Plasmid (m): sc-37878-SH, SSRP1 shRNA (h) Lentiviral Particles: sc-37877-V and SSRP1 shRNA (m) Lentiviral Particles: sc-37878-V.


Molecular Weight of SSRP1: 81 kDa.

Positive Controls: SSRP1 (m): 293T Lysate: sc-127593 or K-562 nuclear extract: sc-2130.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



SSRP1 (D-15): sc-5909. Western blot analysis of SSRP1 expression in non-transfected 293T: sc-117752 (A), mouse SSRP1 transfected 293T: sc-127593 (B) and HeLa (C) whole cell lysates.



SSRP1 (D-15): sc-5909. Western blot analysis of SSRP1 expression in K-562 nuclear extract.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



Try **SSRP1 (D-7): sc-74536** or **SSRP1 (3E4): sc-56782**, our highly recommended monoclonal alternatives to SSRP1 (D-15).