



MHC class I α (KSK 001-03): sc-59311

BACKGROUND

Major histocompatibility complex (MHC) molecules, also designated human leukocyte antigen (HLA) molecules, are cell-surface receptors that bind foreign peptides and present them to T lymphocytes. MHC class I molecules consist of two polypeptide chains, an α or heavy chain, and β -2-Microglobulin, a non-covalently associated protein. Cytotoxic T lymphocytes bind antigenic peptides presented by MHC class I molecules. Antigens that bind to MHC class I molecules are typically 8-10 residues in length and are stabilized in a peptide binding groove. MHC class II molecules are encoded by polymorphic MHC genes and consist of a non-covalent complex of an α and β chain. Helper T lymphocytes bind antigenic peptides presented by MHC class II molecules. MHC class II molecules bind 13-18 amino acid antigenic peptides. Accumulating in endosomal/lysosomal compartments and on the surface of B cells, HLA-DM and -DO molecules regulate binding of exogenous peptides to class II molecules (HLA-DR) by sustaining a conformation that favors peptide exchange. The differential structural properties of MHC class I and class II molecules account for their respective roles in activating different populations of T lymphocytes.

REFERENCES

1. Murphy, D.B., Lo, D., Rath, S., Brinster, R.L., Flavell, R.A., Slanetz, A. and Janeway, C.A. 1989. A novel MHC class II epitope expressed in thymic medulla but not cortex. *Nature* 338: 765-768.
2. Rudensky A.Yu, Rath, S., Preston-Hurlburt, P., Murphy, D.B. and Janeway, C.A. 1991. On the complexity of self. *Nature* 353: 660-662.
3. Agger, R., Petersen, M.S., Toldbod, H.E., Holtz, S., Dagnaes-Hansen, F., Johnsen, B.W., Bolund, L. and Hokland, M. 2000. Characterization of murine dendritic cells derived from adherent blood mononuclear cells *in vitro*. *Scand. J. Immunol.* 52: 138-147.
4. Viret, C. and Janeway, C.A. 2000. Functional and phenotypic evidence for presentation of E α 52-68 structurally related self-peptide(s) in I-E α -deficient mice. *J. Immunol.* 164: 4627-4634.
5. Fischer, G.F. and Mayr, W.R. 2001. Molecular genetics of the HLA complex. *Wien. Klin. Wochenschr.* 113: 814-824.
6. Günther, E. and Walter, L. 2001. The major histocompatibility complex of the rat (*Rattus norvegicus*). *Immunogenetics* 53: 520-542.
7. Little, A.M. and Parham, P. 2001. Polymorphism and evolution of HLA class I and II genes and molecules. *Rev. Immunogenet.* 1: 105-123.
8. Van Kaer, L. 2001. Accessory proteins that control the assembly of MHC molecules with peptides. *Immunol. Res.* 23: 205-214.
9. Zaliauskiene, L., Kang, S., Sparks, K., Zinn, K.R., Schwiebert, L.M., Weaver, C.T. and Collawn, J.F. 2002. Enhancement of MHC class II-restricted responses by receptor-mediated uptake of peptide antigens. *J. Immunol.* 169: 2337-2345.

CHROMOSOMAL LOCATION

Genetic locus: HLA-B (human) mapping to 6p21.3.

SOURCE

MHC class I α (KSK 001-03) is a mouse monoclonal antibody raised against white blood cells of chicken origin.

PRODUCT

Each vial contains 100 μ g IgG₁ in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

MHC class I α (KSK 001-03) is recommended for detection of both the native as well as the denatured forms of MHC class I α of chicken origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1–2 μ g per 100–500 μ g of total protein (1 ml of cell lysate)] and flow cytometry (1 μ g per 1×10^6 cells).

Molecular Weight of MHC class I α : 46 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-mouse IgG-HRP: sc-2005 (dilution range: 1:2000-1:32,000) or Cruz Marker™ compatible goat anti-mouse IgG-HRP: sc-2031 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml).

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.