



Thomsen-Friedenreich (A78-G/A7): sc-59394

BACKGROUND

During metastasis, the ability of malignant cells to form multicellular aggregates via homotypic or heterotypic aggregation and their adhesion to the endothelium are critical. The tumor-associated carbohydrate Thomsen-Friedenreich antigen (Gal-GalNAc) is involved in tumor cell adhesion and tissue invasion. It also causes an immune response, and overexpression of the antigen causes cancer cells to be more sensitive to natural killer cell lysis. The Thomsen-Friedenreich antigen is suppressed in normal healthy cells and represents one of the few chemically well-defined antigens associated with tumor malignancy. The presence of the Thomsen-Friedenreich antigen on the surface of cancer cells may result from a divergence from the normal pathway for O-linked glycosylation in these cells, most likely caused by inappropriate localization of the enzymes involved in synthesis of the disaccharide.

REFERENCES

- Hull, S.R. and Carraway, K.L. 1988. Mechanism of expression of Thomsen-Friedenreich (T) antigen at the cell surface of a mammary adenocarcinoma. *FASEB J.* 2: 2380-2384.
- Glinsky, V.V., Huflejt, M.E., Glinsky, G.V., Deutscher, S.L. and Quinn, T.P. 2000. Effects of Thomsen-Friedenreich antigen-specific peptide P-30 on β -galactoside-mediated homotypic aggregation and adhesion to the endothelium of MDA-MB-435 human breast carcinoma cells. *Cancer Res.* 60: 2584-2588.
- Iraozqui, F.J. and Nores, G.A. 2003. Thomsen-Friedenreich disaccharide immunogenicity. *Curr. Cancer Drug Targets* 3: 433-443.
- Goletz, S., Cao, Y., Danielczyk, A., Ravn, P., Schoeber, U. and Karsten, U. 2004. Thomsen-Friedenreich antigen: the "hidden" tumor antigen. *Adv. Exp. Med. Biol.* 535: 147-162.
- Sotiriadis, J., Shin, S.C., Yim, D., Sieber, D. and Kim, Y.B. 2004. Thomsen-Friedenreich (T) antigen expression increases sensitivity of natural killer cell lysis of cancer cells. *Int. J. Cancer* 111: 388-397.
- Awad, L., Riedner, J. and Vogel, P. 2005. C-linked disaccharide analogue of the Thomsen-Friedenreich (T)-epitope α -O-conjugated to L-serine. *Chemistry* 11: 3565-3573.
- Iraozqui, F.J., Sendra, V.G., Lardone, R.D. and Nores, G.A. 2005. Immune response to Thomsen-Friedenreich disaccharide and glycan engineering. *Immunol. Cell Biol.* 83: 405-412.
- Jeschke, U., Karsten, U., Wiest, I., Schulze, S., Kuhn, C., Friese, K. and Walzel, H. 2006. Binding of galectin-1 (gal-1) to the Thomsen-Friedenreich (TF) antigen on trophoblast cells and inhibition of proliferation of trophoblast tumor cells *in vitro* by gal-1 or an anti-TF antibody. *Histochem. Cell Biol.* 126: 437-444.
- Thors, C., Jansson, B., Helin, H. and Linder, E. 2006. Thomsen-Friedenreich oncofetal antigen in *Schistosoma mansoni*: localization and immunogenicity in experimental mouse infection. *Parasitology* 132: 73-81.

CHROMOSOMAL LOCATION

Genetic locus: TF (human) mapping to 3q22.1.

SOURCE

Thomsen-Friedenreich (A78-G/A7) is a mouse monoclonal antibody raised against neuraminidase treated red blood cells of human origin.

PRODUCT

Each vial contains 50 μ g IgM in 0.5 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

Thomsen-Friedenreich (A78-G/A7) is recommended for detection of Thomsen-Friedenreich of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence and immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.