



Blood Group H2 (0.BG.6): sc-59466

BACKGROUND

Blood-group antigens are generally defined as molecules formed by sequential addition of saccharides to the carbohydrate side chains of lipids and proteins detected on erythrocytes and certain epithelial cells. The A, B and H antigens are reported to undergo modulation during malignant cellular transformation. Blood group related antigens are usually mucin-type and are detected on erythrocytes, certain epithelial cells and in secretions of certain individuals. Sixteen genetically and biosynthetically distinct but inter-related specificities belong to this group of antigens, including A (1 and 2), B, H (1 and 2), M, N, Lewis A, Lewis B, Lewis X, Lewis Y and precursor type 1 chain antigens. The expressions of the H1 and H2 in different cell types are controlled by different genes. Blood Group H1 (O) is a member of a group of murine monoclonal antibodies that detects blood group specificities of the ABH and Lewis systems.

REFERENCES

- Lloyd, K.O., Larson, G., Strömberg, N., Thurin, J. and Karlsson, K.A. 1983. Mouse monoclonal antibody F-3 recognizes the difucosyl type-2 blood group structure. *Immunogenetics* 17: 537-541.
- Anger, B.R., Lloyd, K.O., Oettgen, H.F. and Old, L.J. 1984. Mouse monoclonal IgM antibody against human lung cancer line SK-LC-3 with specificity for H(O) blood group antigen. *Hybridoma* 1: 139-147.
- Sakamoto, J., Yin, B.W. and Lloyd, K.O. 1985. Analysis of the expression of H, Lewis, X, Y and precursor blood group determinants in saliva and red. *Mol. Immunol.* 21: 1093-1098.
- Widmalm, G. and Venable, R.M. 1994. Molecular dynamics simulation and NMR study of a Blood Group H trisaccharide. *Biopolymers* 34: 1079-1088.
- Kojima, H., Nakamura, K., Mineta-Kitajima, R., Sone, Y. and Tamai, Y. 1997. Biosynthesis of a Blood Group H1 antigen by α 1,2-fucosyltransferase in PC12 cells. *Glycoconj. J.* 13: 445-452.
- Yu, L.C., Yang, Y.H., Broadberry, R.E., Chen, Y.H. and Lin, M. 1997. Heterogeneity of the human H blood group α (1,2)fucosyltransferase gene among para-Bombay individuals. *Vox Sang.* 72: 36-40.
- Liu, Y.H., Fujitani, N., Koda, Y. and Kimura, H. 2000. Distribution of H type 1 and of H type 2 antigens of ABO blood group in different cells of human submandibular gland. *J. Histochem. Cytochem.* 46: 69-76.
- Fujitani, N., Liu, Y., Toda, S., Shirouzu, K., Okamura, T. and Kimura, H. 2001. Expression of H type 1 antigen of ABO histo-blood group in normal colon and aberrant expressions of H type 2 and H type 3/4 antigens in colon cancer. *Glycoconj. J.* 17: 331-338.
- Sakuma, Y., Kubushiro, K., Tsukazaki, K. and Nozawa, S. 2003. Involvement of H type 1 carbohydrate antigen in cell adhesion to vascular endothelial cells of human endometrial cancer. *Anticancer Res.* 23: 1277-1282.

SOURCE

Blood Group H2 (0.BG.6) is a mouse monoclonal antibody raised against HEL erythroleukemia cells of human origin.

PRODUCT

Each vial contains 200 μ g IgG₁ in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

Blood Group H2 (0.BG.6) is recommended for detection of Blood Group H2 of human origin by immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500); non cross-reactive with Oh (Bombay) and may cross-react weakly with A1 erythrocytes.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Immunofluorescence: use goat anti-mouse IgG-FITC: sc-2010 (dilution range: 1:100-1:400) or goat anti-mouse IgG-TR: sc-2781 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.