

CD15 (VIM-C6): sc-59533

BACKGROUND

Fucosyltransferases (FucTs) catalyze the covalent association of fucose to different positional linkages on sugar acceptor molecules. The carbohydrate moieties that are generated are covalently attached to cell surfaces and are necessary to ensure a surface contour that satisfies a variety of physiological roles. CD15, also known as Lewis X or Le^X, is a carbohydrate antigen that is generated by FucT-IV (α 1,3-fucosyltransferase IV). Commonly found on the surface of leukocytes and some tumor cells, CD15 is a trisaccharide that is synthesized when FucT-IV transfers an α -fucose residue onto the β -GlcNAc moiety of cellular N-acetylglucosamines. CD15 functions as an adhesion molecule capable of calcium-mediated homotypic binding. Cells with high surface expression of CD15, therefore, exhibit strong self-aggregation (based on CD15-CD15 interaction) in the presence of calcium. Additionally, CD15 is thought to be a ligand for Selectins (proteins involved in mediating leukocyte-specific cellular interactions), further supporting its role as a cell-adhesion protein.

REFERENCES

1. Eggens, I., et al. 1989. Specific interaction between Le^X and Lex determinants. A possible basis for cell recognition in preimplantation embryos and in embryonal carcinoma cells. *J. Biol. Chem.* 264: 9476-9484.
2. Hakomori, S. 1992. Le^X and related structures as adhesion molecules. *Histochem. J.* 24: 771-776.
3. Nimgaonkar, M., et al. 1996. A combination of CD34 selection and complement-mediated immunopurging (anti-CD15 monoclonal antibody) eliminates tumor cells while sparing normal progenitor cells. *J. Hematother.* 5: 39-48.
4. Warren, H.S., et al. 1996. A carbohydrate structure associated with CD15 (Lewis X) on myeloid cells is a novel ligand for human CD2. *J. Immunol.* 156: 2866-2873.
5. Cao, Z., et al. 2001. Role of the Lewis X glycan determinant in corneal epithelial cell adhesion and differentiation. *J. Biol. Chem.* 276: 21714-21723.
6. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 104230. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
7. Iwamori, M., et al. 2007. Characteristic expression of Lewis-antigenic glycolipids in human ovarian carcinoma-derived cells with anticancer drug-resistance. *J. Biochem.* 141: 309-317.

CHROMOSOMAL LOCATION

Genetic locus: FUT4 (human) mapping to 11q21.

SOURCE

CD15 (VIM-C6) is a mouse monoclonal antibody raised against CD15 of human origin.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PRODUCT

Each vial contains 200 μ g IgM kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS


CD15 (VIM-C6) is recommended for detection of CD15 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and flow cytometry (1 μ g per 1×10^6 cells).

Positive Controls: THP-1 cell lysate: sc-2238 or human PBL whole cell lysate.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



CD15 (VIM-C6): sc-59533. Western blot analysis of CD15 expression in human PBL whole cell lysate.

SELECT PRODUCT CITATIONS

1. Wang, C., et al. 2017. Dickkopf-related protein 2 is epigenetically inactivated and suppresses colorectal cancer growth and tumor metastasis by antagonizing Wnt/ β -catenin signaling. *Cell. Physiol. Biochem.* 41: 1709-1724.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.



See **CD15 (C3D-1): sc-19648** for CD15 antibody conjugates, including AC, HRP, FITC, PE, Alexa Fluor® 488, 546, 594, 647, 680 and 790.