



MRP5 (M5II-54): sc-59617

BACKGROUND

The two members of the large family of ABC transporters known to confer multidrug resistance in human cancer cells are the Mdr-1 P-glycoprotein and the multidrug-resistance protein MRP1. MRP1 is an integral membrane protein that contains an MDR-like core, an N-terminal membrane-bound region and a cytoplasmic linker, and it is expressed in various cerebral cells, as well as in lung, testis and peripheral blood. The MRP gene family also includes MRP2, which is alternatively designated cMOAT (for canalicular multispecific organic anion transporter), and MRP3, which are both conjugate export pumps expressed predominantly in hepatocytes. MRP2 localizes exclusively to the apical membrane and is constitutively expressed at a high level in normal liver cells. Conversely, MRP3 localizes to the basolateral membrane where it also mediates the transport of the organic anion S-(2,4-dinitrophenyl)- glutathione toward the basolateral side of the membrane. MRP3 is normally expressed at comparatively lower levels than MRP2 and increases only when secretion across the apical membrane by MRP2 is impaired. MRP6 protein is highly expressed in liver and kidney, whereas MRP4 and MRP5 are detected in various tissues yet at much lower levels of expression.

REFERENCES

1. Versantvoort, C.H., et al. 1995. Regulation by glutathione of drug transport in multidrug-resistant human lung tumour cell lines overexpressing multidrug resistance-associated protein. *Br. J. Cancer* 72: 82-89.
2. Keppler, D. and Konig, J. 1997. Hepatic canalicular membrane 5: expression and localization of the conjugate export pump encoded by the MRP2 (cMRP/cMOAT) gene in liver. *FASEB J.* 11: 509-516.
3. Kool, M., et al. 1997. Analysis of expression of cMOAT (MRP2), MRP3, MRP4, and MRP5, homologues of the multidrug resistance-associated protein gene (MRP1), in human cancer cell lines. *Cancer Res.* 57: 3537-3547.
4. Bakos, E., et al. 1998. Functional multidrug resistance protein (MRP1) lacking the N-terminal transmembrane domain. *J. Biol. Chem.* 273: 32167-32175.
5. Ortiz, D.F., et al. 1999. MRP3, a new ATP-binding cassette protein localized to the canalicular domain of the hepatocyte. *Am. J. Physiol.* 276: 1493-1500.
6. Konig, J., et al. 1999. Characterization of the human multidrug resistance protein isoform MRP3 localized to the basolateral hepatocyte membrane. *Hepatology* 29: 1156-1163.
7. Stockel, B., et al. 2000. Characterization of the 5'-flanking region of the human multidrug resistance protein 2 (MRP2) gene and its regulation in comparison with the multidrug resistance protein 3 (MRP3) gene. *Eur. J. Biochem.* 267: 1347-1358.

CHROMOSOMAL LOCATION

Genetic locus: MRP5 (human) mapping to 3q27.

STORAGE

For immediate and continuous use, store at 4° C for up to one month. For sporadic use, freeze in working aliquots in order to avoid repeated freeze/thaw cycles. If turbidity is evident upon prolonged storage, clarify solution by centrifugation.

SOURCE

MRP5 (M5II-54) is a rat monoclonal antibody raised against amino acids 722-910 of MRP5 of human origin.

PRODUCT

Each vial contains 500 µl culture supernatant containing IgG with < 0.1% sodium azide and 0.7% BSA.

APPLICATIONS

MRP5 (M5II-54) is recommended for detection of MRP5 of human origin by Western Blotting (starting dilution to be determined by researcher, dilution range 1:10-1:200), immunofluorescence (starting dilution to be determined by researcher, dilution range 1:10-1:200) and flow cytometry [1 µg (approximately 10-20 µl) per 1 x 10⁶ cells].

Suitable for use as control antibody for MRP5 siRNA (h): sc-35965, MRP5 shRNA Plasmid (h): sc-35965-SH and MRP5 shRNA (h) Lentiviral Particles: sc-35965-V.

Molecular Weight of MRP5: 185 kDa.

Positive Controls: A549 cell lysate: sc-2413, SK-N-SH cell lysate: sc-2410 or HeLa whole cell lysate: sc-2200.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rat IgG-HRP: sc-2006 (dilution range: 1:2000-1:32,000) or Cruz Marker™ compatible goat anti-rat IgG-HRP: sc-2032 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use goat anti-rat IgG-FITC: sc-2011 (dilution range: 1:100-1:400) or goat anti-rat IgG-TR: sc-2782 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.