

CtIP (K-19): sc-5969

BACKGROUND

CtBP1 is a 48 kDa cellular phosphoprotein that associates with various proteins and functions as a corepressor of transcription. CtBP1 and the related protein CtBP2 are characterized as C-terminal binding protein of adenovirus E1A, and they preferentially associate with the E1A via a 5-amino acid motif, PLDLS, to repress E1A induced oncogenesis and cellular transformation. CtBP1 is expressed from embryo to adult, but CtBP2 is mainly expressed during embryogenesis. During skeletal and T-cell development, CtBP1 and CtBP2 associate with the PLDLS domain of δ EF1, a cellular zinc finger-homeodomain protein, and thereby enhance δ EF1 induced transcriptional silencing. In addition, CtBP complexes with CtIP, a 125 kDa protein that recognizes distinctly different protein motifs from CtBP. CtIP binds to the BRCT repeats within the breast cancer gene BRCA1 and enables CtBP to influence BRCA1 activity. CtIP/CtBP binding to BRCA1 inhibits the transactivation of the p21 promoter, and it is critical for regulating p21 transcription in response to DNA damage.

REFERENCES

- Sollerbrant, K., et al. 1996. The CtBP binding domain in the adenovirus E1A protein controls CR1-dependent transactivation. *Nucleic Acids Res.* 24: 2578-2584.
- Sekido, R., et al. 1997. Two mechanisms in the action of repressor δ EF1: binding site competition with an activator and active repression. *Genes Cells* 2: 771-783.
- Schaeper, U., et al. 1998. Interaction between a cellular protein that binds to the C-terminal region of adenovirus E1A (CtBP) and a novel cellular protein is disrupted by E1A through a conserved PLDLS motif. *J. Biol. Chem.* 273: 8549-8552.
- Turner, J., et al. 1998. Cloning and characterization of mCtBP2, a corepressor that associates with basic Krüppel-like factor and other mammalian transcriptional regulators. *EMBO J.* 17: 5129-5140.
- Yu, X., et al. 1998. The C-terminal (BRCT) domains of BRCA1 interact *in vivo* with CtIP, a protein implicated in the CtBP pathway of transcriptional repression. *J. Biol. Chem.* 273: 25388-25392.

CHROMOSOMAL LOCATION

Genetic locus: RBBP8 (human) mapping to 18q11.2; Rbbp8 (mouse) mapping to 18 A1.

SOURCE

CtIP (K-19) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of CtIP of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-5969 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

CtIP (K-19) is recommended for detection of CtIP of mouse and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).


CtIP (K-19) is also recommended for detection of CtIP in additional species, including equine, canine, porcine and avian.

Suitable for use as control antibody for CtIP siRNA (h): sc-37765, CtIP siRNA (m): sc-37766, CtIP shRNA Plasmid (h): sc-37765-SH, CtIP shRNA Plasmid (m): sc-37766-SH, CtIP shRNA (h) Lentiviral Particles: sc-37765-V and CtIP shRNA (m) Lentiviral Particles: sc-37766-V.

Molecular Weight of CtIP: 125 kDa.

Positive Controls: T24 cell lysate: sc-2292, Jurkat nuclear extract: sc-2132 or CtIP (m): 293T Lysate: sc-119500.

DATA



CtIP (K-19): sc-5969. Western blot analysis of CtIP expression in non-transfected: sc-117752 (A) and mouse CtIP transfected: sc-119500 (B) 293T whole cell lysates.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.


 MONOS
Satisfation
Guaranteed

Try **CtIP (D-4): sc-271339** or **CtIP (F-2): sc-28324**, our highly recommended monoclonal alternatives to CtIP (K-19).