



SP-D (IIE11): sc-59698

CKGROUND

Pulmonary surfactant is primarily responsible for lowering the surface tension at the air-liquid interface in the alveoli, a process that is essential for normal respiration. Pulmonary surfactant is a mixture of phospholipids and proteins, including four distinct surfactant-associated proteins (SPs), SP-A, SP-B, SP-C and SP-D. SP-B and SP-C are predominantly hydrophobic proteins that associate with lipids to promote the absorption of surfactant phospholipids and to reduce the surface tension in the alveoli. SP-A and SP-D are large multimeric proteins belonging to the family of calcium-dependent lectins, designated collectins, which contribute to the innate immune system. Both SP-A and SP-D have been shown to protect against microbial challenge through binding to the lipid components of the bacterial cell wall and facilitating the rapid removal of microbials.

REFERENCES

1. Glasser, S.W., Korfhagen, T.R., Bruno, M.D., Dey, C. and Whitsett, J.A. 1990. Structure and expression of the pulmonary surfactant protein SP-C gene in the mouse. *J. Biol. Chem.* 265: 21986-21991.
2. Hawgood, S. and Shiffer, K. 1991. Structures and properties of the surfactant-associated proteins. *Annu. Rev. Physiol.* 53: 375-394.
3. Johansson, J., Jornvall, H. and Curstedt, T. 1992. Human surfactant polypeptide SP-B. Disulfide bridges, C-terminal end and peptide analysis of the airway form. *FEBS Lett.* 301: 165-167.
4. Crouch, E., Rust, K., Veile, R., Donis-Keller, H. and Grosso, L. 1993. Genomic organization of human surfactant protein D (SP-D). SP-D is encoded on chromosome 10q22.2-23.1. *J. Biol. Chem.* 268: 2976-2983.
5. Rooney, S.A., Young, S.L. and Mendelson, C.R. 1994. Molecular and cellular processing of lung surfactant. *FASEB J.* 8: 957-967.
6. Johansson, J. and Curstedt, T. 1997. Molecular structures and interactions of pulmonary surfactant components. *Eur. J. Biochem.* 244: 675-693.
7. Reid, K.B. 1998. Functional roles of the lung surfactant proteins SP-A and SP-D in innate immunity. *Immunobiology* 199: 200-207.
8. Wert, S.E., Yoshida, M., LeVine, A.M., Ikegami, M., Jones, T., Ross, G.F., Fisher, J.H., Korfhagen, T.R. and Whitsett, J.A. 2000. Increased metalloproteinase activity, oxidant production and emphysema in SP-D gene-inactivated mice. *Proc. Natl. Acad. Sci. USA* 97: 5972-5977.
9. McCormack, F.X. and Whitsett, J.A. 2002. The pulmonary collectins, SP-A and SP-D, orchestrate innate immunity in the lung. *J. Clin. Invest.* 109: 707-712.

CHROMOSOMAL LOCATION

Genetic locus: SFTPD (human) mapping to 10q22.3; Sftpd (mouse) mapping to 14 B.

SOURCE

SP-D (IIE11) is a mouse monoclonal antibody raised against full length native SP-D of rat origin.

PRODUCT

Each vial contains 50 µg IgG_{2b} in 500 µl PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

SP-D (IIE11) is recommended for detection of SP-D of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with SP-A.

Suitable for use as control antibody for SP-D siRNA (h): sc-36541, SP-D siRNA (m): sc-36542, SP-D shRNA Plasmid (h): sc-36541-SH, SP-D shRNA Plasmid (m): sc-36542-SH, SP-D shRNA (h) Lentiviral Particles: sc-36541-V and SP-D shRNA (m) Lentiviral Particles: sc-36542-V.

Molecular Weight of SP-D: 43 kDa.

Positive Controls: rat lung extract: sc-2396 or human lung extract: sc-363767.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.